


# Dyonic maximal supergravity from massive IIA and its Chern-Simons duals

Adolfo Guarino  
(Nikhef, Amsterdam)

December 1st , Uppsala



**Nikhef**




With Daniel Jafferis and Oscar Varela :

[arXiv:1504.08009](https://arxiv.org/abs/1504.08009) , [arXiv:1508.04432](https://arxiv.org/abs/1508.04432) , [arXiv:1509.02526](https://arxiv.org/abs/1509.02526)


# Outlook




Motivation




Deformed  $SO(8)$ -gauged supergravity



Deformed  $ISO(7)$ -gauged supergravity



Massive type IIA strings on  $S^6$



New  $AdS_4$   $N=2$  solution in massive IIA and its  $CFT_3$  dual



Motivation

# electric-magnetic deformations

- The uniqueness of the maximal ( $N=8$ ) supergravities is historically inherited from their connection to sphere reductions

$$\text{AdS}_5 \times S^5 \text{ (D3-brane)} \quad , \quad \text{AdS}_4 \times S^7 \text{ (M2-brane)} \quad , \quad \text{AdS}_7 \times S^4 \text{ (M5-brane)}$$

- $N=8$  supergravity in 4D admits a **deformation parameter**  $c$  yielding **inequivalent** theories. It is an **electric/magnetic** deformation

$$D = \partial - g (A^{\text{elec}} - c \tilde{A}_{\text{mag}})$$

$g$  = 4D gauge coupling  
 $c$  = **deformation param.**

- There are two generic situations :
  - 1) Family of  $\text{SO}(8)_c$  theories :  $c = [0, \sqrt{2} - 1]$  is a continuous param. [ similar for  $\text{SO}(p,q)_c$  ]
  - 2) Family of  $\text{ISO}(7)_c$  theories :  $c = 0 \text{ or } 1$  is an (on/off) param. [ same for  $\text{ISO}(p,q)_c$  ]

[ Dall'Agata, Inverso, Trigiante '12 ]

[ Dall'Agata, Inverso, Marrani '14 ]

The questions arise:

- Does such an electric / magnetic deformation of 4D maximal supergravity enjoy a string / M-theory origin, or is it just a 4D feature ?


Obstruction for  $SO(8)_c$  , *cf.* [ Lee, Strickland-Constable & Waldram '15 ]

[ de Wit & Nicolai '13 ]

- For deformed 4D supergravities with supersymmetric  $AdS_4$  vacua, are these  $AdS_4$  /  $CFT_3$ -dual to any identifiable 3d CFT ?

# A new 10D / 4D / 3d correspondence

*massive* IIA on  $S^6$   $\ll$  ISO(7)<sub>c</sub>-gauged sugra  $\gg$  SU(N)<sub>k</sub> C-S-M theory



$g_c$  = elec/mag deformation in 4D


$\hat{F}_{(0)}$  = Romans mass in 10D

$k$  = Chern-Simons level in 3d

[ Schwarz '04 ]

[ AG, Jafferis, Varela '15 ]

[ AG, Varela '15 ]



Deformed  $SO(8)$ -gauged supergravity

# $N = 8$ supergravities in 4D

- SUGRA :    metric + 8 gravitini + 28 vectors + 56 dilatini + 70 scalars  
                  (s = 2)            (s = 3/2)            (s = 1)            (s = 1/2)            (s = 0)

Ungauged (abelian) supergravity: Reduction of M-theory on a *torus*  $T^7$   
down to 4D produces  $N = 8$  supergravity with  $G = U(1)^{28}$  [ Cremmer & Julia '79 ]

Gauged (non-abelian) supergravity: Reduction of M-theory on a *sphere*  $S^7$   
down to 4D produces  $N = 8$  supergravity with  $G = SO(8)$  [ de Wit & Nicolai '82 ]

\*  $SO(8)$ -gauged supergravity believed to be **unique** for 30 years...

... but ... is this true?



# Framework to study $N = 8$ supergravities in 4D

[ de Wit, Samtleben & Trigiante '03 , '07 ]

Gauging procedure : Part of the **global  $E_7$  symmetry** group is promoted to a local symmetry group  $G$  (gauging)

[  $\alpha = 1, \dots, 133$  ]

**Embedding tensor** : It is a “selector” specifying which **generators of  $E_7$**  (there are 133!!) become the gauge symmetry  $G$  and, therefore, have associated gauge fields.

Formulation in terms of **56** vectors  $A_\mu^M$ , though...  $M = 1, \dots, 56 = 28$  (elec) + **28** (mag)

*Sp(56) Elec/Mag group*

$$A_\mu = A_\mu^M \Theta_M^\alpha t_\alpha$$

*Redundancy!!*

$$X_M = \Theta_M^\alpha t_\alpha \quad \Rightarrow \quad [X_M, X_N] = X_{MN}^P X_P \quad \text{with} \quad X_{MN}^P = \Theta_M^\alpha [t_\alpha]_N^P$$

\* Closure of the gauge algebra :  $\Omega^{MN} \Theta_M^\alpha \Theta_N^\beta = 0$

*Only 28 physical l.c. of vectors!!*

# A family of $G = \text{SO}(8)$ supergravities in 4D


- Choose  $G = \text{SO}(8)$
- Solve  $\Omega^{MN} \Theta_M^\alpha \Theta_N^\beta = 0 \Rightarrow$  One-parameter ( $c$ ) family of  $\text{SO}(8)_c$  theories !!

[ Dall'Agata, Inverso & Trigiante '12 ]

- Immediate questions :

- |  |                              |
|--|------------------------------|
| 1) What?                                   | (Yes, surprising but true)   |
| 2) Are these $c$ -theories equivalent?     | (No)                         |
| 3) Are there new $\text{AdS}_4$ solutions? | (Yes)                        |
| 4) Higher-dimensional origin?              | (Good question... )          |
| 5) $\text{AdS}_4/\text{CFT}_3$ dual?       | (Good question too... ABJ? ) |

Physical meaning in 4D : electric/magnetic deformation



$$D = \partial - g (A^{\text{elec}} - c \tilde{A}_{\text{mag}})$$

Physical meaning 10D / 11D ...



Holographic  $\text{AdS}_4/\text{CFT}_3$  meaning ...



In this talk we are going to investigate the electric/magnetic deformation of a different N=8 supergravity closely related to the  $G = SO(8)$  theory ...

... the  $G = ISO(7) = SO(7) \ltimes \mathbb{R}^7$  supergravity !!

electric / magnetic  
deformation



higher-dimensional  
origin



Holographic  
AdS<sub>4</sub>/CFT<sub>3</sub> dual ?





Deformed ISO(7)-gauged supergravity

# A family of $G = \text{ISO}(7)$ supergravities in 4D

- Choose  $G = \text{ISO}(7)$
- Solve  $\Omega^{MN} \Theta_M^\alpha \Theta_N^\beta = 0 \Rightarrow$  One-parameter ( $c$ ) family of  $\text{ISO}(7)_c$  theories !!

[ Hull '84 (electric) ]


[ Dall'Agata, Inverso, Marrani '14 ]

- Immediate questions :

- |  |                             |
|--|-----------------------------|
| 1) What?                                   | (Yes, and still surprising) |
| 2) Are these $c$ -theories equivalent?     | (No)                        |
| 3) Are there new $\text{AdS}_4$ solutions? | (Yes)                       |
| 4) Higher-dimensional origin?              | (Yes)                       |
| 5) $\text{AdS}_4/\text{CFT}_3$ dual?       | (Yes)                       |



# Physical meaning in 4D = electric/magnetic deformation



$$D = \partial - g A_{\text{SO}(7)}^{\text{elec}} - g (A_{\mathbb{R}^7}^{\text{elec}} - c \tilde{A}_{\mathbb{R}^7 \text{ mag}})$$

# Deformed ISO(7)<sub>c</sub> Lagrangian ( $m = g\mathbf{c}$ )

$$M = 1, \dots, 56$$

$$\Lambda = 1, \dots, 28$$

$$I = 1, \dots, 7$$

$$\begin{aligned} \mathcal{L}_{\text{bos}} = & (R - V) \text{vol}_4 - \frac{1}{48} D\mathcal{M}_{\text{MIN}} \wedge *D\mathcal{M}^{\text{MIN}} + \frac{1}{2} \mathcal{I}_{\Lambda\Sigma} \mathcal{H}_{(2)}^\Lambda \wedge *\mathcal{H}_{(2)}^\Sigma - \frac{1}{2} \mathcal{R}_{\Lambda\Sigma} \mathcal{H}_{(2)}^\Lambda \wedge \mathcal{H}_{(2)}^\Sigma \\ & + m \left[ \mathcal{B}^I \wedge (\tilde{\mathcal{H}}_{(2)I} - \frac{g}{2} \delta_{IJ} \mathcal{B}^J) - \frac{1}{4} \tilde{\mathcal{A}}_I \wedge \tilde{\mathcal{A}}_J \wedge (d\mathcal{A}^{IJ} + \frac{g}{2} \delta_{KL} \mathcal{A}^{IK} \wedge \mathcal{A}^{JL}) \right] \end{aligned}$$

◆ Setting  $m = 0$ , all the magnetic pieces in the Lagrangian disappear.

## \* Ingredients :

- Electric vectors (21 + 7):  $\mathcal{A}^{IJ} = \mathcal{A}^{[IJ]}$  [SO(7)] and  $\mathcal{A}^I$  [R<sup>7</sup>] with  $\mathcal{H}_{(2)}^\Lambda = (\mathcal{H}_{(2)}^{IJ}, \mathcal{H}_{(2)}^I)$
- Auxiliary magnetic vectors (7):  $\tilde{\mathcal{A}}_I$  [R<sup>7</sup>] with  $\tilde{\mathcal{H}}_{(2)I}$  field strength
- E<sub>7</sub>/SU(8) scalars :  $\mathcal{M}_{\text{MIN}}$
- Auxiliary two-forms (7):  $\mathcal{B}^I$  [R<sup>7</sup>]
- Topological term :  $m$  [ ... ]
- Scalar potential :  $V(\mathcal{M}) = \frac{g^2}{672} X_{\text{MN}}^{\text{R}} X_{\text{PQ}}^{\text{S}} \mathcal{M}^{\text{MP}} (\mathcal{M}^{\text{NQ}} \mathcal{M}_{\text{RS}} + 7 \delta_{\text{R}}^{\text{Q}} \delta_{\text{S}}^{\text{N}})$

# A truncation : $G_0 = \text{SU}(3)$ invariant sector

[ Warner '83 ]

- **Truncation** : Retaining the fields and couplings which are invariant (singlets) under the action of a subgroup  $G_0 \subset \text{ISO}(7)$

- **SU(8) R-symmetry branching** : **gravitini**  $\mathbf{8} \rightarrow \mathbf{1} + \mathbf{1} + \mathbf{3} + \bar{\mathbf{3}} \Rightarrow \mathcal{N} = 2$  SUSY

- **Scalars fields** :  $\mathbf{70} \rightarrow \mathbf{1} (\times 6) + \text{non-singlets} \Rightarrow 6$  real scalars  $(\varphi, \chi; \phi, a, \zeta, \tilde{\zeta})$

- **Vector fields** :  $\mathbf{56} \rightarrow \mathbf{1} (\times 4) + \text{non-singlets} \Rightarrow$  vectors  $(A^0, A^1; \tilde{A}_0, \tilde{A}_1)$

- $N = 2$  gauged supergravity with  $G = \text{U}(1) \times \text{SO}(1,1)_c$  coupled to **1 vector & 1 hyper**

$$\mathcal{M}_{\text{scalar}} = \frac{\text{SU}(1,1)}{\text{U}(1)} \times \frac{\text{SU}(2,1)}{\text{U}(2)}$$

- **Scalar potential** : 
$$V = \frac{1}{2} g^2 \left[ e^{4\phi-3\varphi} (1 + e^{2\varphi} \chi^2)^3 - 12 e^{2\phi-\varphi} (1 + e^{2\varphi} \chi^2) - 12 e^{2\phi+\varphi} \rho^2 (1 - 3 e^{2\varphi} \chi^2) \right. \\ \left. - 24 e^\varphi + 12 e^{4\phi+\varphi} \chi^2 \rho^2 (1 + e^{2\varphi} \chi^2) + 12 e^{4\phi+\varphi} \rho^4 (1 + 3 e^{2\varphi} \chi^2) \right] \\ - \frac{1}{2} gm \chi e^{4\phi+3\varphi} (12 \rho^2 + 2\chi^2) + \frac{1}{2} m^2 e^{4\phi+3\varphi} ,$$

*AdS critical points !!*

# AdS<sub>4</sub> solutions

| $\mathcal{N}$     | $G_0$     | $c^{-1/3} \chi$      | $c^{-1/3} e^{-\varphi}$           | $c^{-1/3} \rho$        | $c^{-1/3} e^{-\phi}$              | $\frac{1}{4} g^{-2} c^{1/3} V_0$    | $M^2 L^2$                                       |
|-------------------|-----------|----------------------|-----------------------------------|------------------------|-----------------------------------|-------------------------------------|---|
| $\mathcal{N} = 1$ | $G_2$     | $-\frac{1}{2^{7/3}}$ | $\frac{5^{1/2} 3^{1/2}}{2^{7/3}}$ | $-\frac{1}{2^{7/3}}$   | $\frac{5^{1/2} 3^{1/2}}{2^{7/3}}$ | $-\frac{2^{22/3} 3^{1/2}}{5^{5/2}}$ | $4 \pm \sqrt{6}, -\frac{1}{6}(11 \pm \sqrt{6})$ |
| $\mathcal{N} = 2$ | $U(3)$    | $-\frac{1}{2}$       | $\frac{3^{1/2}}{2}$               | 0                      | $\frac{1}{2^{1/2}}$               | $-3^{3/2}$                          | $3 \pm \sqrt{17}, 2, 2$                         |
| $\mathcal{N} = 1$ | $SU(3)$   | $\frac{1}{2^2}$      | $\frac{3^{1/2} 5^{1/2}}{2^2}$     | $-\frac{3^{1/2}}{2^2}$ | $\frac{5^{1/2}}{2^2}$             | $-\frac{2^6 3^{3/2}}{5^{5/2}}$      | $4 \pm \sqrt{6}, 4 \pm \sqrt{6}$                |
| $\mathcal{N} = 0$ | $SO(6)_+$ | 0                    | $2^{1/6}$                         | 0                      | $\frac{1}{2^{5/6}}$               | $-3 2^{5/6}$                        | $6, 6, -\frac{3}{4}, 0$                         |
| $\mathcal{N} = 0$ | $SO(7)_+$ | 0                    | $\frac{1}{5^{1/6}}$               | 0                      | $\frac{1}{5^{1/6}}$               | $-\frac{3 5^{7/6}}{2^2}$            | $6, -\frac{12}{5}, -\frac{6}{5}, -\frac{6}{5}$  |
| $\mathcal{N} = 0$ | $G_2$     | $\frac{1}{2^{4/3}}$  | $\frac{3^{1/2}}{2^{4/3}}$         | $\frac{1}{2^{4/3}}$    | $\frac{3^{1/2}}{2^{4/3}}$         | $-\frac{2^{10/3}}{3^{1/2}}$         | $6, 6, -1, -1$                                  |
| $\mathcal{N} = 0$ | $SU(3)$   | 0.455                | 0.838                             | 0.335                  | 0.601                             | -5.864                              | 6.214, 5.925, 1.145, -1.284                     |
| $\mathcal{N} = 0$ | $SU(3)$   | 0.270                | 0.733                             | 0.491                  | 0.662                             | -5.853                              | 6.230, 5.905, 1.130, -1.264                     |

◆ Relevant for holographic RG flows (in progress...)

# The truncated Lagrangian and its dual formulation

- The Lagrangian contains a **non-dynamical tensor field  $B^0$**  :

$$\begin{aligned}
 \mathcal{L} = & (R - V) \text{vol}_4 + \frac{3}{2} [d\varphi \wedge *d\varphi + e^{2\varphi} d\chi \wedge *d\chi] \\
 & + 2 d\phi \wedge *d\phi + \frac{1}{2} e^{2\phi} [D\zeta \wedge *D\zeta + D\tilde{\zeta} \wedge *D\tilde{\zeta}] \\
 & + \frac{1}{2} e^{4\phi} [Da + \frac{1}{2}(\zeta D\tilde{\zeta} - \tilde{\zeta} D\zeta)] \wedge *[Da + \frac{1}{2}(\zeta D\tilde{\zeta} - \tilde{\zeta} D\zeta)] \\
 & + \frac{1}{2} \mathcal{I}_{\Lambda\Sigma} H_{(2)}^\Lambda \wedge *H_{(2)}^\Sigma - \frac{1}{2} \mathcal{R}_{\Lambda\Sigma} H_{(2)}^\Lambda \wedge H_{(2)}^\Sigma + m B^0 \wedge d\tilde{A}_0 + \frac{1}{2} g m B^0 \wedge B^0
 \end{aligned}$$

- Solving the EOM for the magnetic vector, one finds

$$H_{(3)}^0 = e^{4\phi} * (Da + \frac{1}{2} (\zeta D\tilde{\zeta} - \tilde{\zeta} D\zeta)) \quad \longrightarrow \quad \text{scalar-tensor duality!!}$$

- The dual Lagrangian contains a **dynamical tensor field  $H^0 = dB^0$**  :

$$\begin{aligned}
 \tilde{\mathcal{L}} = & (R - V) \text{vol}_4 + \frac{1}{2} e^{-4\phi} H_{(3)}^0 \wedge *H_{(3)}^0 + \frac{3}{2} [d\varphi \wedge *d\varphi + e^{2\varphi} d\chi \wedge *d\chi] \\
 & + 2 d\phi \wedge *d\phi + \frac{1}{2} e^{2\phi} [D\zeta \wedge *D\zeta + D\tilde{\zeta} \wedge *D\tilde{\zeta}] \\
 & + \frac{1}{2} \mathcal{I}_{\Lambda\Sigma} H_{(2)}^\Lambda \wedge *H_{(2)}^\Sigma - \frac{1}{2} \mathcal{R}_{\Lambda\Sigma} H_{(2)}^\Lambda \wedge H_{(2)}^\Sigma \\
 & - H_{(3)}^0 \wedge \left[ g A^0 + \frac{1}{2} (\zeta D\tilde{\zeta} - \tilde{\zeta} D\zeta) \right] + \frac{1}{2} g m B^0 \wedge B^0 .
 \end{aligned}$$

- ◆ Natural duality frame to investigate possible higher-dimensional origin!!

[ N=2 Calabi-Yau and coset reds w/ fluxes ]

[ Polchinski & Strominger '95 ]

[ Louis & Micu '02 ]

[ Kashani-Poor '07 ]

[ Kashani-Poor & Cassani '09 ]

Dual formulations seem to be crucial  
to understand  
higher-dimensional origins!!

... let's give up the Lagrangian.

# Tensor hierarchy

[ de Wit, Nicolai & Samtleben '08 ]

[ Bergshoeff, Hartong, Hohm, Huebscher & Ortin '09 ]

- **Idea:** To describe the dynamics of a 4D theory in terms of a set of  $p$ -form fields with  $p = 0, 1, 2, 3$  [no Lagrangian!]

- Restricted  $SL(7)$ -covariant field content [ index  $I$  ]

|                     |               |   |              |
|---------------------|---------------|---|--------------|
| $21' + 7' + 21 + 7$ | vectors :     | $\mathcal{A}^{IJ}, \mathcal{A}^I, \tilde{\mathcal{A}}_{IJ}, \tilde{\mathcal{A}}_I,$ |              |
| $48 + 7'$           | two-forms :   | $\mathcal{B}_I^J, \mathcal{B}^I,$   | [out of 133] |
| $28'$               | three-forms : | $\mathcal{C}^{IJ},$   | [out of 912] |

- Two-form field strengths [  $21' + 7' + 21 + 7$  ]

$$\mathcal{H}_{(2)}^{IJ} = d\mathcal{A}^{IJ} - g \delta_{KL} \mathcal{A}^{IK} \wedge \mathcal{A}^{LJ},$$

$$\mathcal{H}_{(2)}^I = d\mathcal{A}^I - g \delta_{JK} \mathcal{A}^{IJ} \wedge \mathcal{A}^K + \frac{1}{2} m \mathcal{A}^{IJ} \wedge \tilde{\mathcal{A}}_J + m \mathcal{B}^I,$$

$$\tilde{\mathcal{H}}_{(2)IJ} = d\tilde{\mathcal{A}}_{IJ} + g \delta_{K[I} \mathcal{A}^{KL} \wedge \tilde{\mathcal{A}}_{J]L} + g \delta_{K[I} \mathcal{A}^K \wedge \tilde{\mathcal{A}}_{J]} - m \tilde{\mathcal{A}}_I \wedge \tilde{\mathcal{A}}_J + 2g \delta_{K[I} \mathcal{B}_{J]}^K,$$

$$\tilde{\mathcal{H}}_{(2)I} = d\tilde{\mathcal{A}}_I - \frac{1}{2} g \delta_{IJ} \mathcal{A}^{JK} \wedge \tilde{\mathcal{A}}_K + g \delta_{IJ} \mathcal{B}^J,$$

- Three-form field strengths [ **48 + 7'** ]

$$\begin{aligned}
\mathcal{H}_{(3)I}{}^J &= DB_I{}^J + \frac{1}{2}\mathcal{A}^{JK} \wedge d\tilde{\mathcal{A}}_{IK} + \frac{1}{2}\mathcal{A}^J \wedge d\tilde{\mathcal{A}}_I + \frac{1}{2}\tilde{\mathcal{A}}_{IK} \wedge d\mathcal{A}^{JK} + \frac{1}{2}\tilde{\mathcal{A}}_I \wedge d\mathcal{A}^J \\
&\quad - \frac{1}{2}g\delta_{KL}\mathcal{A}^{JK} \wedge \mathcal{A}^{LM} \wedge \tilde{\mathcal{A}}_{IM} - \frac{1}{2}g\delta_{KL}\mathcal{A}^{JK} \wedge \mathcal{A}^L \wedge \tilde{\mathcal{A}}_I \\
&\quad + \frac{1}{6}g\delta_{IK}\mathcal{A}^{JL} \wedge \mathcal{A}^{KM} \wedge \tilde{\mathcal{A}}_{LM} - \frac{1}{3}g\delta_{IK}\mathcal{A}^{(J} \wedge \mathcal{A}^{K)L} \wedge \tilde{\mathcal{A}}_L \\
&\quad - \frac{1}{2}m\mathcal{A}^{JK} \wedge \tilde{\mathcal{A}}_I \wedge \tilde{\mathcal{A}}_K - 2g\delta_{IK}\mathcal{C}^{JK} - \frac{1}{7}\delta_I^J(\text{trace}) ,
\end{aligned}$$

$$\mathcal{H}_{(3)}^I = DB^I - \frac{1}{2}\mathcal{A}^{IJ} \wedge d\tilde{\mathcal{A}}_J - \frac{1}{2}\tilde{\mathcal{A}}_J \wedge d\mathcal{A}^{IJ} + \frac{1}{2}g\delta_{JK}\mathcal{A}^{IJ} \wedge \mathcal{A}^{KL} \wedge \tilde{\mathcal{A}}_L ,$$

- Four-form field strengths [ **28'** ]

$$\begin{aligned}
\mathcal{H}_{(4)}^{IJ} &= DC^{IJ} - \mathcal{H}_{(2)}^{K(I} \wedge \mathcal{B}_K{}^{J)} + \mathcal{H}_{(2)}^{(I} \wedge \mathcal{B}^{J)} - \frac{1}{2}m\mathcal{B}^I \wedge \mathcal{B}^J - \frac{1}{6}\mathcal{A}^{K(I} \wedge \tilde{\mathcal{A}}_{KL} \wedge d\mathcal{A}^{J)L} \\
&\quad + \frac{1}{6}\mathcal{A}^{IK} \wedge \mathcal{A}^{JL} \wedge d\tilde{\mathcal{A}}_{KL} - \frac{1}{6}\mathcal{A}^{K(I} \wedge \tilde{\mathcal{A}}_K \wedge d\mathcal{A}^{J)} - \frac{1}{3}\mathcal{A}^{K(I} \wedge \mathcal{A}^{J)} \wedge d\tilde{\mathcal{A}}_K \\
&\quad - \frac{1}{6}\mathcal{A}^{(I} \wedge \tilde{\mathcal{A}}_K \wedge d\mathcal{A}^{J)K} - \frac{1}{6}g\delta_{KL}\mathcal{A}^{K(I} \wedge \mathcal{A}^{J)M} \wedge \mathcal{A}^{LN} \wedge \tilde{\mathcal{A}}_{MN} \\
&\quad + \frac{1}{6}g\delta_{KL}\mathcal{A}^{K(I} \wedge \mathcal{A}^{J)} \wedge \mathcal{A}^{LM} \wedge \tilde{\mathcal{A}}_M - \frac{1}{6}g\delta_{KL}\mathcal{A}^{K(I} \wedge \mathcal{A}^{J)M} \wedge \mathcal{A}^L \wedge \tilde{\mathcal{A}}_M \\
&\quad - \frac{1}{8}m\mathcal{A}^{IK} \wedge \mathcal{A}^{JL} \wedge \tilde{\mathcal{A}}_K \wedge \tilde{\mathcal{A}}_L .
\end{aligned}$$



# Consistency checks

- Closed set of Bianchi identities

$$\begin{aligned}
 D\mathcal{H}_{(2)}^{IJ} &= 0 \quad , \quad D\mathcal{H}_{(2)}^I = m \mathcal{H}_{(3)}^I \quad , \quad D\tilde{\mathcal{H}}_{(2)IJ} = -2g \mathcal{H}_{(3)[I}{}^K \delta_{J]K} \quad , \quad D\tilde{\mathcal{H}}_{(2)I} = g \delta_{IJ} \mathcal{H}_{(3)}^J \quad , \\
 D\mathcal{H}_{(3)I}{}^J &= \mathcal{H}_{(2)}^{JK} \wedge \tilde{\mathcal{H}}_{(2)IK} + \mathcal{H}_{(2)}^J \wedge \tilde{\mathcal{H}}_{(2)I} - 2g \delta_{IK} \mathcal{H}_{(4)}^{JK} - \frac{1}{7} \delta_I^J \text{ (trace)} \quad , \\
 D\mathcal{H}_{(3)}^I &= -\mathcal{H}_{(2)}^{IJ} \wedge \tilde{\mathcal{H}}_{(2)J} \quad , \quad D\mathcal{H}_{(4)}^{IJ} \equiv 0 \quad .
 \end{aligned}$$

- Closed set of duality relations [right number of d.o.f] [short-hand notation]

$$\begin{aligned}
 \tilde{\mathcal{H}}_{(2)IJ} &= -\frac{1}{2} \mathcal{I}_{[IJ][KL]} * \mathcal{H}_{(2)}^{KL} - \mathcal{I}_{[IJ][K8]} * \mathcal{H}_{(2)}^K + \frac{1}{2} \mathcal{R}_{[IJ][KL]} \mathcal{H}_{(2)}^{KL} + \mathcal{R}_{[IJ][K8]} \mathcal{H}_{(2)}^K \quad , \\
 \tilde{\mathcal{H}}_{(2)I} &= -\frac{1}{2} \mathcal{I}_{[I8][KL]} * \mathcal{H}_{(2)}^{KL} - \mathcal{I}_{[I8][K8]} * \mathcal{H}_{(2)}^K + \frac{1}{2} \mathcal{R}_{[I8][KL]} \mathcal{H}_{(2)}^{KL} + \mathcal{R}_{[I8][K8]} \mathcal{H}_{(2)}^K \quad , \\
 \mathcal{H}_{(3)I}{}^J &= -\frac{1}{12} (t_I^J)_{\mathbb{M}^{\mathbb{P}}} \mathcal{M}_{\mathbb{NP}} * D\mathcal{M}^{\mathbb{MN}} - \frac{1}{7} \delta_I^J \text{ (trace)} \quad , \\
 \mathcal{H}_{(3)}^I &= -\frac{1}{12} (t_8^I)_{\mathbb{M}^{\mathbb{P}}} \mathcal{M}_{\mathbb{NP}} * D\mathcal{M}^{\mathbb{MN}} \quad , \\
 \mathcal{H}_{(4)}^{IJ} &= \frac{1}{84} X_{\mathbb{NQ}}^{\mathbb{S}} \left( (t_K^{(I|})_{\mathbb{P}^{\mathbb{R}}} \mathcal{M}^{|J)KN} + (t_8^{(I|})_{\mathbb{P}^{\mathbb{R}}} \mathcal{M}^{|J)8N} \right) (\mathcal{M}^{\mathbb{PQ}} \mathcal{M}_{\mathbb{RS}} + 7 \delta_{\mathbb{S}}^{\mathbb{P}} \delta_{\mathbb{R}}^{\mathbb{Q}}) \text{vol}_4 \quad .
 \end{aligned}$$


[  $t$ 's are  $\text{SL}(7) \times \mathbb{R}^7$  generators ]

- Closed set of SUSY transformations

**Q** : Why to bother with all these tensor hierarchy issues?

**A** : Because the tensor hierarchy allows us to derive simple formulas to embed the *4D dynamics* into a higher-dimensional theory

Remember : *No index, no clue.* Good luck trying to embed  $V$  into higher dimensions...



Massive type IIA strings on  $S^6$

# Collecting clues

- The deformed  $\text{ISO}(7)_c$  gauging has its  $\text{SO}(7)$  piece untouched by the deformation. This points towards an undeformed  $S^6$  description in higher dimension.

- If the higher-dimensional geometry is not affected, it should then be the higher-dimensional theory the one changing. The massive IIA theory by Romans proves a natural candidate.

[ Romans '86 ]

- The Romans mass parameter  $\hat{F}_{(0)}$  is a discrete (on/off) deformation, exactly as the parameter  $c$  in the deformed  $\text{ISO}(7)_c$  theory.

- The  $\text{SU}(3)$ -invariant sector of the  $\text{ISO}(7)_c$  theory connects to  $\text{CY}_3$  reductions of massive IIA upon dualisation of some field. It is then natural to believe that the embedding of the full  $\text{ISO}(7)_c$  theory would require an enlarged set of duality relations... tensor hierarchy!!

# Derivation of the IIA embedding [ 4-step process ]

\* Step 1 : 10D KK decomposition that leaves 4D spacetime symmetry manifest

$A(x,y)$ 's and  $B(x,y)$ 's fields

\* Step 2 : Redefinitions of the  $A$ 's and  $B$ 's fields to conform 4D SUSY transformations

$C(x,y)$ 's fields

\* Step 3 : Connection to actual 4D fields by dressing up with  $S^6$  geometrical data

$$C(x, y) = \text{geometry}(y) \times \mathcal{C}(x)$$

\* Step 4 : Plug and play

\* **Step 1** : 10D redefinitions ( KK decomp) that leave 4D spacetime symmetry manifest

$$\text{SO}(1, 9) \rightarrow \text{SO}(1, 3) \times \text{SO}(6)$$

Then one has :

$$d\hat{s}_{10}^2 = \Delta^{-1} ds_4^2 + g_{mn} (dy^m + B^m)(dy^n + B^n) ,$$

$$\begin{aligned} \hat{A}_{(3)} &= \frac{1}{6} A_{\mu\nu\rho} dx^\mu \wedge dx^\nu \wedge dx^\rho + \frac{1}{2} A_{\mu\nu m} dx^\mu \wedge dx^\nu \wedge (dy^m + B^m) \\ &+ \frac{1}{2} A_{\mu mn} dx^\mu \wedge (dy^m + B^m) \wedge (dy^n + B^n) \\ &+ \frac{1}{6} A_{mnp} (dy^m + B^m) \wedge (dy^n + B^n) \wedge (dy^p + B^p) , \end{aligned}$$

$$\hat{B}_{(2)} = \frac{1}{2} B_{\mu\nu} dx^\mu \wedge dx^\nu + B_{\mu m} dx^\mu \wedge (dy^m + B^m) + \frac{1}{2} B_{mn} (dy^m + B^m) \wedge (dy^n + B^n) ,$$

$$\hat{A}_{(1)} = A_\mu dx^\mu + A_m (dy^m + B^m) ,$$

In terms of representations of  $\text{SL}(6)$  [ index  $m$  ] :

|  |                             |              |  |
|--|-----------------------------|--------------|--|
|  | <b>1</b>                    | metric :     | $ds_4^2$ ,   |
|  | <b>21 + 6 + 1 + 20 + 15</b> | scalars :    | $g_{mn}$ , $A_m$ , $\hat{\phi}$ , $A_{mnp}$ , $B_{mn}$ , |
|  | <b>6' + 1 + 15 + 6</b>      | vectors :    | $B_\mu^m$ , $A_\mu$ , $A_{\mu mn}$ , $B_{\mu m}$ ,       |
|  | <b>6 + 1</b>                | two-forms :  | $A_{\mu\nu m}$ , $B_{\mu\nu}$ ,                          |
|  | <b>1</b>                    | three-form : | $A_{\mu\nu\rho}$ .                                       |

$$\Delta^2 \equiv \frac{\det g_{mn}}{\det \hat{g}_{mn}}$$

\* Step 2 : Non-linear field redefinitions to conform 4D SUSY transformations

- Vectors :  $C_\mu^{m8} \equiv B_\mu^m$  ,  $C_\mu^{78} \equiv A_\mu$  ,  $\tilde{C}_{\mu mn} \equiv A_{\mu mn} - A_\mu B_{mn}$  ,  $\tilde{C}_{\mu m7} \equiv B_{\mu m}$
- Two-forms :  $C_{\mu\nu m}^8 \equiv -A_{\mu\nu m} + C_{[\mu}^{n8} \tilde{C}_{\nu]nm} + C_{[\mu}^{78} \tilde{C}_{\nu]m7}$  ,  $C_{\mu\nu 7}^8 \equiv -B_{\mu\nu} + C_{[\mu}^{m8} \tilde{C}_{\nu]m7}$
- Three-form :  $C_{\mu\nu\rho}^{88} \equiv A_{\mu\nu\rho} - C_{[\mu}^{m8} C_{\nu}^{n8} \tilde{C}_{\rho]mn} + C_{[\mu}^{m8} C_{\nu}^{78} \tilde{C}_{\rho]m7} + 3 C_{[\mu}^{78} C_{\nu\rho]7}^8$

These can be rearranged into representations of SL(7) [ index  $I$  ]

$$C_\mu^{I8} = (C_\mu^{m8}, C_\mu^{78}) \quad \tilde{C}_{\mu IJ} = (\tilde{C}_{\mu mn}, \tilde{C}_{\mu m7}) \quad C_{\mu\nu I}^8 = (C_{\mu\nu m}^8, C_{\mu\nu 7}^8) \quad C_{\mu\nu\rho}^{88}$$

with 10D SUSY transfs:

$$\delta C_\mu^{I8} = i V^{I8}{}_{AB} \left( \bar{\epsilon}^A \psi_\mu^B + \frac{1}{2\sqrt{2}} \bar{\epsilon}_C \gamma_\mu \chi^{ABC} \right) + \text{h.c.} ,$$

$$\delta \tilde{C}_{\mu IJ} = -i V_{IJ}{}_{AB} \left( \bar{\epsilon}^A \psi_\mu^B + \frac{1}{2\sqrt{2}} \bar{\epsilon}_C \gamma_\mu \chi^{ABC} \right) + \text{h.c.} ,$$

...

Mimicking the 4D tensor hierarchy !!

The result is then a set of  $SL(7)$ -covariant 10D fields :

$$\begin{array}{lll}
\mathbf{1} & \text{metric :} & ds_4^2(x, y) , \\
\mathbf{7}' + \mathbf{21} & \text{generalised vielbeine :} & V^{I8}_{AB}(x, y) , \tilde{V}_{IJAB}(x, y) , \\
\mathbf{7}' + \mathbf{21} & \text{vectors :} & C_\mu^{I8}(x, y) , \tilde{C}_{\mu IJ}(x, y) , \\
\mathbf{7} & \text{two-forms :} & C_{\mu\nu I}{}^8(x, y) , \\
\mathbf{1} & \text{three-form :} & C_{\mu\nu\rho}{}^{88}(x, y) .
\end{array}$$

that is to be connected with the  $SL(7)$ -covariant 4D fields of the tensor hierarchy :

$$\begin{array}{lll}
\mathbf{1} & \text{metric :} & ds_4^2(x) , \\
\mathbf{21}' + \mathbf{7}' + \mathbf{21} + \mathbf{7} & \text{coset representatives :} & \mathcal{V}^{IJij}(x) , \mathcal{V}^{I8ij}(x) , \tilde{\mathcal{V}}_{IJ}{}^{ij}(x) , \tilde{\mathcal{V}}_{I8}{}^{ij}(x) , \\
\mathbf{21}' + \mathbf{7}' + \mathbf{21} + \mathbf{7} & \text{vectors :} & \mathcal{A}_\mu^{IJ}(x) , \mathcal{A}_\mu^I(x) , \tilde{\mathcal{A}}_{\mu IJ}(x) , \tilde{\mathcal{A}}_{\mu I}(x) , \\
\mathbf{48} + \mathbf{7}' & \text{two-forms :} & \mathcal{B}_{\mu\nu I}{}^J(x) , \mathcal{B}_{\mu\nu}{}^I(x) , \\
\mathbf{28}' & \text{three-forms :} & \mathcal{C}_{\mu\nu\rho}{}^{IJ}(x) ,
\end{array}$$

> This connection is established by using geometrical data of the  $S^6$  !!



\* **Step 3** : Connecting 4D [SL(7)] and 10D [SL(6)] fields using the  $S^6$  geometrical data in a “dressing up” process

[ vectors ]

$$C_{\mu}{}^{m8}(x, y) = \frac{1}{2} g K_{IJ}^m(y) \mathcal{A}_{\mu}{}^{IJ}(x) \quad , \quad C_{\mu}{}^{78}(x, y) = -\mu_I(y) \mathcal{A}_{\mu}{}^I(x) \quad ,$$

$$\tilde{C}_{\mu mn}(x, y) = \frac{1}{4} K_{mn}^{IJ}(y) \tilde{\mathcal{A}}_{\mu IJ}(x) \quad , \quad \tilde{C}_{\mu m7}(x, y) = -g^{-1} (\partial_m \mu^I)(y) \tilde{\mathcal{A}}_{\mu I}(x)$$

[ two-forms ]

$$C_{\mu\nu m}{}^8(x, y) = -g^{-1} (\mu_I \partial_m \mu^J)(y) \mathcal{B}_{\mu\nu J}{}^I(x)$$

$$C_{\mu\nu 7}{}^8(x, y) = \mu_I(y) \mathcal{B}_{\mu\nu}{}^I(x)$$

[ three-form ]

$$C_{\mu\nu\rho}{}^{88}(x, y) = (\mu_I \mu_J)(y) \mathcal{C}_{\mu\nu\rho}{}^{IJ}(x)$$

◆  $S^6$  geometrical data : embedding coordinates  $\mu^I$ , Killing vectors  $K_{IJ}^m$  and tensors  $K_{IJ}^{mn}$

\* Step 4 : Plug and play... so that the final embedding of  $\text{ISO}(7)_c$  into type IIA is

$$d\hat{s}_{10}^2 = \Delta^{-1} ds_4^2 + g_{mn} Dy^m Dy^n ,$$

$$\begin{aligned} \hat{A}_{(3)} = & \mu_I \mu_J (C^{IJ} + \mathcal{A}^I \wedge \mathcal{B}^J + \frac{1}{6} \mathcal{A}^{IK} \wedge \mathcal{A}^{JL} \wedge \tilde{\mathcal{A}}_{KL} + \frac{1}{6} \mathcal{A}^I \wedge \mathcal{A}^{JK} \wedge \tilde{\mathcal{A}}_K) \\ & + g^{-1} (\mathcal{B}_J^I + \frac{1}{2} \mathcal{A}^{IK} \wedge \tilde{\mathcal{A}}_{KJ} + \frac{1}{2} \mathcal{A}^I \wedge \tilde{\mathcal{A}}_J) \wedge \mu_I D\mu^J + \frac{1}{2} g^{-2} \tilde{\mathcal{A}}_{IJ} \wedge D\mu^I \wedge D\mu^J \\ & - \frac{1}{2} \mu_I B_{mn} \mathcal{A}^I \wedge Dy^m \wedge Dy^n + \frac{1}{6} A_{mnp} Dy^m \wedge Dy^n \wedge Dy^p , \end{aligned}$$

$$\hat{B}_{(2)} = -\mu_I (\mathcal{B}^I + \frac{1}{2} \mathcal{A}^{IJ} \wedge \tilde{\mathcal{A}}_J) - g^{-1} \tilde{\mathcal{A}}_I \wedge D\mu^I + \frac{1}{2} B_{mn} Dy^m \wedge Dy^n ,$$

$$\hat{A}_{(1)} = -\mu_I \mathcal{A}^I + A_m Dy^m . \quad ($$

where we have defined :  $Dy^m \equiv dy^m + \frac{1}{2} g K_{IJ}^m \mathcal{A}^{IJ}$  ,  $D\mu^I \equiv d\mu^I - g \mathcal{A}^{IJ} \mu_J$

The scalars are embedded as

$$\begin{aligned} g^{mn} &= \frac{1}{4} g^2 \Delta \mathcal{M}^{IJKL} K_{IJ}^m K_{KL}^n , & B_{mn} &= -\frac{1}{2} \Delta g_{mp} K_{IJ}^p \partial_n \mu^K \mathcal{M}^{IJ}_{K8} , \\ A_m &= \frac{1}{2} g \Delta g_{mn} K_{IJ}^n \mu_K \mathcal{M}^{IJK8} , & A_{mnp} &= \frac{1}{8} g \Delta g_{mq} K_{IJ}^q K_{np}^{KL} \mathcal{M}^{IJ}_{KL} + A_m B_{np} . \end{aligned}$$

# Remarks and consistency checks

- 10D (bosonic) SUSY transformations exactly reduce to those of the 4D tensor hierarchy.
- Computing the 10D field strengths  $\hat{F}_{(2)} = d\hat{A}_{(1)} + m\hat{B}_{(2)}$ , etc. one finds

$$\hat{F}_{(4)} = \mu_I \mu_J \mathcal{H}_{(4)}^{IJ} + g^{-1} \mathcal{H}_{(3)J}^I \wedge \mu_I D\mu^J + \frac{1}{2} g^{-2} \tilde{\mathcal{H}}_{(2)IJ} \wedge D\mu^I \wedge D\mu^J + \dots, \quad [\text{FR parameter}]$$

$$\hat{H}_{(3)} = -\mu_I \mathcal{H}_{(3)}^I - g^{-1} \tilde{\mathcal{H}}_{(2)I} \wedge D\mu^I + \dots,$$

$$\hat{F}_{(2)} = -\mu_I \mathcal{H}_{(2)}^I + g^{-1} (g \delta_{IJ} \mathcal{A}^J - m \tilde{\mathcal{A}}_I) \wedge D\mu^I + \dots,$$

which are expressed in terms of the 4D tensor hierarchy. The parameter  $m=gc$  only appears through standard Romans' redefinitions of  $\hat{F}_{(p)}$  in 10D (formulas also valid for massless IIA)

[Hull & Warner '88 (electric)]


- The set of Bianchi identities of the above 10D field strengths reduces to

$$D\mathcal{H}_{(2)}^{IJ} = 0, \quad D\mathcal{H}_{(2)}^I = m \mathcal{H}_{(3)}^I, \quad D\tilde{\mathcal{H}}_{(2)IJ} = -2g \mathcal{H}_{(3)[I}^K \delta_{J]K}, \quad D\tilde{\mathcal{H}}_{(2)I} = g \delta_{IJ} \mathcal{H}_{(3)}^J,$$

$$D\mathcal{H}_{(3)I}^J = \mathcal{H}_{(2)}^{JK} \wedge \tilde{\mathcal{H}}_{(2)IK} + \mathcal{H}_{(2)}^J \wedge \tilde{\mathcal{H}}_{(2)I} - 2g \delta_{IK} \mathcal{H}_{(4)}^{JK} - \frac{1}{7} \delta_I^J (\text{trace}),$$

$$D\mathcal{H}_{(3)}^I = -\mathcal{H}_{(2)}^{IJ} \wedge \tilde{\mathcal{H}}_{(2)J}, \quad D\mathcal{H}_{(4)}^{IJ} \equiv 0.$$

which exactly matches the one of the 4D tensor hierarchy.



New  $\text{AdS}_4$   $N=2$  solution in massive IIA and its  $\text{CFT}_3$  dual

# $N=2$ solution of massive type IIA

[ Luest & Tsimpis '09 ]  
[ see also Petrini & Zaffaroni '09 ]

- Using the uplifting formulae, it is a straightforward exercise to obtain the 10D embedding of a 4D critical point. An example is the  $N=2 \& U(3)$   $AdS_4$  point of the  $ISO(7)_c$  theory

$$d\hat{s}_{10}^2 = L^2 \frac{(3 + \cos 2\alpha)^{\frac{1}{2}}}{(5 + \cos 2\alpha)^{-\frac{1}{8}}} \left[ ds^2(AdS_4) + \frac{3}{2} d\alpha^2 + \frac{6 \sin^2 \alpha}{3 + \cos 2\alpha} ds^2(\mathbb{CP}^2) + \frac{9 \sin^2 \alpha}{5 + \cos 2\alpha} \boldsymbol{\eta}^2 \right],$$

$$e^{\hat{\phi}} = e^{\phi_0} \frac{(5 + \cos 2\alpha)^{3/4}}{3 + \cos 2\alpha}, \quad \hat{H}_{(3)} = 24\sqrt{2} L^2 e^{\frac{1}{2}\phi_0} \frac{\sin^3 \alpha}{(3 + \cos 2\alpha)^2} \boldsymbol{J} \wedge d\alpha,$$

$$L^{-1} e^{\frac{3}{4}\phi_0} \hat{F}_{(2)} = -4\sqrt{6} \frac{\sin^2 \alpha \cos \alpha}{(3 + \cos 2\alpha)(5 + \cos 2\alpha)} \boldsymbol{J} - 3\sqrt{6} \frac{(3 - \cos 2\alpha)}{(5 + \cos 2\alpha)^2} \sin \alpha d\alpha \wedge \boldsymbol{\eta},$$

$$L^{-3} e^{\frac{1}{4}\phi_0} \hat{F}_{(4)} = 6 \text{vol}_4 + 12\sqrt{3} \frac{7 + 3 \cos 2\alpha}{(3 + \cos 2\alpha)^2} \sin^4 \alpha \text{vol}_{\mathbb{CP}^2} + 18\sqrt{3} \frac{(9 + \cos 2\alpha) \sin^3 \alpha \cos \alpha}{(3 + \cos 2\alpha)(5 + \cos 2\alpha)} \boldsymbol{J} \wedge d\alpha \wedge \boldsymbol{\eta},$$

where we have introduced the quantities  $L^2 \equiv 2^{-\frac{5}{8}} 3^{-1} g^{-2} c^{\frac{1}{12}}$  and  $e^{\phi_0} \equiv 2^{\frac{1}{4}} c^{-\frac{5}{6}}$

- ◆ The angle  $0 \leq \alpha \leq \pi$  locally foliates  $S^6$  with  $S^5$  regarded as Hopf fibrations over  $\mathbb{CP}^2$

# CFT<sub>3</sub> candidate and matching of free energies

[ Schwarz '04 ]

[ Gaiotto & Tomasiello '09 ]

- We propose an N=2 Chern-Simons-matter theory with simple gauge group SU(N), level k and only adjoint matter, as the CFT of the N=2 massive IIA solution.

- The 3d free energy  $F = -\text{Log}(Z)$ , where Z is the partition function on the CFT on a Euclidean S<sup>3</sup> can be computed via localisation over supersymmetric configurations

[ Pestun '07 ] [ Jafferis '10 ]

[ Jafferis, Klebanov, Pufu & Safdi '11 ]

$$Z = \int \prod_{i=1}^N \frac{d\lambda_i}{2\pi} \prod_{i<j=1}^N \left( 2 \sinh^2 \left( \frac{\lambda_i - \lambda_j}{2} \right) \right) \times \prod_{i,j=1}^N \left( \exp \left( \ell \left( \frac{1}{3} + \frac{i}{2\pi} (\lambda_i - \lambda_j) \right) \right) \right)^3 e^{\frac{ik}{4\pi} \sum \lambda_i^2}$$

where  $\lambda_i$  are the Coulomb branch parameters. In the  $N \gg k$  limit, the result is given by

$$F = \frac{3^{13/6} \pi}{40} \left( \frac{32}{27} \right)^{2/3} k^{1/3} N^{5/3}$$

- The gravitational free energy can be computed from the warp factor in the N=2 massive IIA solution. Using the charge quantisation condition  $N = -(2\pi\ell_s)^{-5} \int_{S^6} e^{\frac{1}{2}\phi} \hat{F}_{(4)} + \hat{B}_{(2)} \wedge d\hat{A}_{(3)} + \frac{1}{6} \hat{F}_{(0)} \hat{B}_{(2)}^3$  for the D2-brane, one finds

$$F = \frac{16\pi^3}{(2\pi\ell_s)^8} \int_{S^6} e^{8A} \text{vol}_6 = \frac{\pi}{5} 2^{1/3} 3^{1/6} k^{1/3} N^{5/3} \quad \text{provided}$$

$$g c = \hat{F}_{(0)} = k / (2\pi\ell_s)$$

# Summary


- 1) We have introduced a new method to embed 4D theories into 10D. Using this method, we have connected the dyonic N=8 supergravity with  $ISO(7)_c$  gauging to massive IIA reductions on  $S^6$ .
- 2) Any 4D configuration (AdS, DWs, BH) is embedded into 10D via the uplifting formulas. As an example, we found a new  $AdS_4 \times S^6$  solution of massive IIA based on an N=2&U(3)  $AdS_4$  vacuum.
- 3) We propose a  $CFT_3$  dual for the N=2  $AdS_4 \times S^6$  solution of massive IIA based on the D2-brane field theory. In the massive IIA case, there is a CS-term and a superpotential that make the theory flowing to a conformal phase (IR). This translates into the appearance of supersymmetric  $AdS_4$  vacua in the deformed  $ISO(7)_c$  supergravity theory.
- 4) For the new N=2 massive IIA solution, the gravitational and FT free energies do match provided

$$g c = \hat{F}_{(0)} = k/(2\pi\ell_s)$$

- 5) Extension to other dimensions? EGG / EFT for massive IIA?

Tack så mycket !!





Extra material

# More AdS<sub>4</sub> critical points

[here] = arXiv:1508.04432

| SUSY              | bos. sym.          | $M^2 L^2$  | stability | ref.         |
|-------------------|--------------------|--|-----------|--------------|
| $\mathcal{N} = 3$ | SO(4)              | $3(1 \pm \sqrt{3})^{(1)}$ , $(1 \pm \sqrt{3})^{(6)}$ , $-\frac{9}{4}^{(4)}$ , $-2^{(18)}$ , $-\frac{5}{4}^{(12)}$ , $0^{(22)}$<br>$(3 \pm \sqrt{3})^{(3)}$ , $\frac{15}{4}^{(4)}$ , $\frac{3}{4}^{(12)}$ , $0^{(6)}$               | yes       | [30]         |
| $\mathcal{N} = 2$ | U(3)               | $(3 \pm \sqrt{17})^{(1)}$ , $-\frac{20}{9}^{(12)}$ , $-2^{(16)}$ , $-\frac{14}{9}^{(18)}$ , $2^{(3)}$ , $0^{(19)}$<br>$4^{(1)}$ , $\frac{28}{9}^{(6)}$ , $\frac{4}{9}^{(12)}$ , $0^{(9)}$  | yes       | [15], [here] |
| $\mathcal{N} = 1$ | G <sub>2</sub>     | $(4 \pm \sqrt{6})^{(1)}$ , $-\frac{1}{6}(11 \pm \sqrt{6})^{(27)}$ , $0^{(14)}$<br>$\frac{1}{2}(3 \pm \sqrt{6})^{(7)}$ , $0^{(14)}$   | yes       | [4]          |
| $\mathcal{N} = 1$ | SU(3)              | $(4 \pm \sqrt{6})^{(2)}$ , $-\frac{20}{9}^{(12)}$ , $-2^{(8)}$ , $-\frac{8}{9}^{(12)}$ , $\frac{7}{9}^{(6)}$ , $0^{(28)}$<br>$6^{(1)}$ , $\frac{28}{9}^{(6)}$ , $\frac{25}{9}^{(6)}$ , $2^{(1)}$ , $\frac{4}{9}^{(6)}$ , $0^{(8)}$ | yes       | [here]       |
| $\mathcal{N} = 0$ | SO(7) <sub>+</sub> | $6^{(1)}$ , $-\frac{12}{5}^{(27)}$ , $-\frac{6}{5}^{(35)}$ , $0^{(7)}$<br>$\frac{12}{5}^{(7)}$ , $0^{(21)}$  | no        | [3]          |
| $\mathcal{N} = 0$ | SO(6) <sub>+</sub> | $6^{(2)}$ , $-3^{(20)}$ , $-\frac{3}{4}^{(20)}$ , $0^{(28)}$<br>$6^{(1)}$ , $\frac{9}{4}^{(12)}$ , $0^{(15)}$  | no        | [3]          |
| $\mathcal{N} = 0$ | G <sub>2</sub>     | $6^{(2)}$ , $-1^{(54)}$ , $0^{(14)}$<br>$3^{(14)}$ , $0^{(14)}$  | yes       | [4]          |
| $\mathcal{N} = 0$ | SU(3)              | see (3.44)<br>see (3.45)   | yes       | [here]       |
| $\mathcal{N} = 0$ | SU(3)              | see (3.46)<br>see (3.47)   | yes       | [here]       |
| $\mathcal{N} = 0$ | SO(4)              | see (5.12)<br>see (5.13)   | yes       | [here]       |

# SUSY transformations of the tensor hierarchy

[ vielbein and scalars ]

$$\begin{aligned}\delta e_\mu^\alpha &= \frac{1}{4} \bar{\epsilon}_i \gamma^\alpha \psi_\mu^i + \frac{1}{4} \bar{\epsilon}^i \gamma^\alpha \psi_{\mu i} \\ \delta \mathcal{V}_M^{ij} &= \frac{1}{\sqrt{2}} \mathcal{V}_M^{kl} \left( \bar{\epsilon}^{[i} \chi^{jkl]} + \frac{1}{4!} \varepsilon^{ijklmnpq} \bar{\epsilon}_m \chi_{npq} \right)\end{aligned}$$

[ vectors ]

$$\begin{aligned}\delta \mathcal{A}_\mu^{IJ} &= i \mathcal{V}^{IJ}{}_{ij} \left( \bar{\epsilon}^i \psi_\mu^j + \frac{1}{2\sqrt{2}} \bar{\epsilon}_k \gamma_\mu \chi^{ijk} \right) + \text{h.c.} \\ \delta \mathcal{A}_\mu^I &= i \mathcal{V}^{I8}{}_{ij} \left( \bar{\epsilon}^i \psi_\mu^j + \frac{1}{2\sqrt{2}} \bar{\epsilon}_k \gamma_\mu \chi^{ijk} \right) + \text{h.c.} \\ \delta \tilde{\mathcal{A}}_{\mu IJ} &= -i \tilde{\mathcal{V}}_{IJ}{}_{ij} \left( \bar{\epsilon}^i \psi_\mu^j + \frac{1}{2\sqrt{2}} \bar{\epsilon}_k \gamma_\mu \chi^{ijk} \right) + \text{h.c.} \\ \delta \tilde{\mathcal{A}}_{\mu I} &= -i \tilde{\mathcal{V}}_{I8}{}_{ij} \left( \bar{\epsilon}^i \psi_\mu^j + \frac{1}{2\sqrt{2}} \bar{\epsilon}_k \gamma_\mu \chi^{ijk} \right) + \text{h.c.}\end{aligned}$$

[ two-forms ]

$$\begin{aligned}\delta \mathcal{B}_{\mu\nu J}^I &= \left[ -\frac{2}{3} (\mathcal{V}^{IK}{}_{jk} \tilde{\mathcal{V}}_{JK}{}^{ik} + \mathcal{V}^{I8}{}_{jk} \tilde{\mathcal{V}}_{J8}{}^{ik} + \tilde{\mathcal{V}}_{JK}{}_{jk} \mathcal{V}^{IK}{}^{ik} + \tilde{\mathcal{V}}_{J8}{}_{jk} \mathcal{V}^{I8}{}^{ik}) \bar{\epsilon}_i \gamma_{[\mu} \psi_{\nu]}^j \right. \\ &\quad \left. - \frac{\sqrt{2}}{3} (\mathcal{V}^{IK}{}_{ij} \tilde{\mathcal{V}}_{JK}{}^{kl} + \mathcal{V}^{I8}{}_{ij} \tilde{\mathcal{V}}_{J8}{}^{kl}) \bar{\epsilon}^{[i} \gamma_{\mu\nu} \chi^{jkl]} + \text{h.c.} \right] \\ &\quad + (\mathcal{A}_{[\mu}^{IK} \delta \tilde{\mathcal{A}}_{\nu]JK} + \mathcal{A}_{[\mu}^I \delta \tilde{\mathcal{A}}_{\nu]J} + \tilde{\mathcal{A}}_{[\mu|JK} \delta \mathcal{A}_{|\nu]}^{IK} + \tilde{\mathcal{A}}_{[\mu|J} \delta \mathcal{A}_{|\nu]}^I) - \frac{1}{7} \delta_J^I (\text{trace}), \\ \delta \mathcal{B}_{\mu\nu}^I &= \left[ \frac{2}{3} (\mathcal{V}^{IJ}{}_{jk} \tilde{\mathcal{V}}_{J8}{}^{ik} + \tilde{\mathcal{V}}_{J8}{}_{jk} \mathcal{V}^{IJ}{}^{ik}) \bar{\epsilon}_i \gamma_{[\mu} \psi_{\nu]}^j + \frac{\sqrt{2}}{3} \mathcal{V}^{IJ}{}_{ij} \tilde{\mathcal{V}}_{J8}{}^{kl} \bar{\epsilon}^{[i} \gamma_{\mu\nu} \chi^{jkl]} + \text{h.c.} \right] \\ &\quad - (\mathcal{A}_{[\mu}^{IJ} \delta \tilde{\mathcal{A}}_{\nu]J} + \tilde{\mathcal{A}}_{[\mu|J} \delta \mathcal{A}_{|\nu]}^{IJ}).\end{aligned}$$

[ three-forms ]

$$\begin{aligned}
\delta\mathcal{C}_{\mu\nu\rho}{}^{IJ} = & \left[ -\frac{4i}{7} \left( \mathcal{V}^{K(I}{}_{jl}(\mathcal{V}^{J)Llk} \tilde{\mathcal{V}}_{KLik} + \tilde{\mathcal{V}}_{KL}{}^{lk} \mathcal{V}^{J)L}{}_{ik} \right) \right. \\
& + \mathcal{V}^{K(I}{}_{jl}(\mathcal{V}^{J)8lk} \tilde{\mathcal{V}}_{K8ik} + \tilde{\mathcal{V}}_{K8}{}^{lk} \mathcal{V}^{J)8}{}_{ik} \\
& + \mathcal{V}^{(I|8}{}_{jl}(\mathcal{V}^{|J)Klk} \tilde{\mathcal{V}}_{K8ik} + \tilde{\mathcal{V}}_{K8}{}^{lk} \mathcal{V}^{|J)K}{}_{ik} \left. \right) \bar{\epsilon}^i \gamma_{[\mu\nu} \psi_{\rho]}^j \\
& + i\frac{\sqrt{2}}{3} \left( \mathcal{V}^{K(I|hi} \mathcal{V}^{|J)L}{}_{[ij]} \tilde{\mathcal{V}}_{KL|kl} + \mathcal{V}^{K(Ihi} \mathcal{V}^{J)8}{}_{[ij]} \tilde{\mathcal{V}}_{K8|kl} \right. \\
& \left. + \mathcal{V}^{(I|8hi} \mathcal{V}^{|J)K}{}_{[ij]} \tilde{\mathcal{V}}_{K8|kl} \right) \bar{\epsilon}_h \gamma_{\mu\nu\rho} \chi^{jkl} + \text{h.c.} \left. \right] \\
& - 3 \left( \mathcal{B}_{[\mu\nu|K}{}^{(I} \delta\mathcal{A}_{|\rho]}^{J)K} + \mathcal{B}_{[\mu\nu}{}^{(I} \delta\mathcal{A}_{\rho]}^{J)} \right) \\
& + \mathcal{A}_{[\mu}{}^{K(I} (\mathcal{A}_{\nu}^{J)L} \delta\tilde{\mathcal{A}}_{\rho]KL} + \tilde{\mathcal{A}}_{\nu KL} \delta\mathcal{A}_{\rho]}^{J)L}) + \mathcal{A}_{[\mu}{}^{K(I} (\mathcal{A}_{\nu}^{J)} \delta\tilde{\mathcal{A}}_{\rho]K} + \tilde{\mathcal{A}}_{\nu K} \delta\mathcal{A}_{\rho]}^{J)}) \\
& + \mathcal{A}_{[\mu}{}^{(I} (\mathcal{A}_{\nu}^{J)K} \delta\tilde{\mathcal{A}}_{\rho]K} + \tilde{\mathcal{A}}_{\nu K} \delta\mathcal{A}_{\rho]}^{J)K}) .
\end{aligned}$$

... all scalars, vectors and the fermions should be kept !!

## Scalar potential and three-form potentials

- How does the scalar potential potential  $V$  fit in the duality hierarchy ?

$$\Theta_{\mathbb{M}}^{\alpha} \mathcal{H}_{(4)\alpha}^{\mathbb{M}} = -2V \text{vol}_4$$

- In our deformed  $\text{ISO}(7)_c$  theory, one has four-form field strengths

$$g \delta_{IJ} \mathcal{H}_{(4)}^{IJ} + m \tilde{\mathcal{H}}_{(4)} = -2V \text{vol}_4 \quad [ \mathbf{28}' + \mathbf{1} \text{ of } \text{SL}(7) ]$$

where we need the **SL(7)-singlet** four-form field strength  $\tilde{\mathcal{H}}_{(4)}$  dual to the magnetic ET

- Consistency requires also the three-form field strength  $\mathcal{H}_{(3)}$  rendering  $\mathcal{H}_{(3)I}{}^J$  traceful

$$\mathcal{H}_{(3)} = \frac{1}{12} (t_8^8)_{\mathbb{M}}^{\mathbb{P}} \mathcal{M}_{\mathbb{NP}} * D\mathcal{M}^{\mathbb{MN}},$$

$$\tilde{\mathcal{H}}_{(4)} = \frac{1}{84} X_{\mathbb{NQ}}^{\mathbb{S}} (t_8^K)_{\mathbb{P}}^{\mathbb{R}} \mathcal{M}_{8K}^{\mathbb{N}} \mathcal{M}^{\mathbb{PQ}} \mathcal{M}_{\mathbb{RS}} \text{vol}_4$$

\* Extended BI's :

$$D\mathcal{H}_{(3)} = \mathcal{H}_{(2)}^{IJ} \wedge \tilde{\mathcal{H}}_{(2)IJ} + \mathcal{H}_{(2)}^I \wedge \tilde{\mathcal{H}}_{(2)I} - 2g \delta_{IJ} \mathcal{H}_{(4)}^{IJ} - 14m \tilde{\mathcal{H}}_{(4)}$$

$$D\tilde{\mathcal{H}}_{(4)} \equiv 0,$$

# Generalised vielbein

$$V^{I8}{}_{AB} = (V^{m8}{}_{AB}, V^{78}{}_{AB}) \quad , \quad \tilde{V}_{IJ}{}_{AB} = (\tilde{V}_{mn}{}_{AB}, \tilde{V}_{m7}{}_{AB}) :$$

with components

$$\begin{aligned} V^{m8}{}_{AB} &= -\frac{1}{4} \Delta^{-\frac{1}{2}} e_a{}^m (\Gamma^a C^{-1})^{AB} , \\ V^{78}{}_{AB} &= -\frac{1}{4} e^{-\frac{3}{4}\hat{\phi}} \Delta^{-\frac{1}{2}} (\Gamma_7 C^{-1})^{AB} - V^{m8}{}_{AB} A_m , \\ \tilde{V}_{m7}{}^{AB} &= \frac{1}{4} e^{\frac{1}{2}\hat{\phi}} \Delta^{-\frac{1}{2}} e_m{}^a (\Gamma_a \Gamma_7 C^{-1})^{AB} + V^{n8}{}_{AB} B_{nm} , \\ \tilde{V}_{mn}{}^{AB} &= \frac{1}{4} e^{-\frac{1}{4}\hat{\phi}} \Delta^{-\frac{1}{2}} e_m{}^a e_n{}^b (\Gamma_{ab} C^{-1})^{AB} + V^{p8}{}_{AB} (A_{pmn} - 2B_{p[m} A_{n]}) \\ &\quad + V^{78}{}_{AB} B_{mn} + 2\tilde{V}_{[m|7}{}^{AB} A_{|n]} \end{aligned}$$

and

$$\begin{aligned} V^{m8}{}_{AB} &= \frac{1}{4} \Delta^{-\frac{1}{2}} e_a{}^m (C\Gamma^a)_{AB} , \\ V^{78}{}_{AB} &= \frac{1}{4} e^{-\frac{3}{4}\hat{\phi}} \Delta^{-\frac{1}{2}} (C\Gamma_7)_{AB} - V^{m8}{}_{AB} A_m , \\ \tilde{V}_{m7}{}_{AB} &= \frac{1}{4} e^{\frac{1}{2}\hat{\phi}} \Delta^{-\frac{1}{2}} e_m{}^a (C\Gamma_a \Gamma_7)_{AB} + V^{n8}{}_{AB} B_{nm} , \\ \tilde{V}_{mn}{}_{AB} &= \frac{1}{4} e^{-\frac{1}{4}\hat{\phi}} \Delta^{-\frac{1}{2}} e_m{}^a e_n{}^b (C\Gamma_{ab})_{AB} + V^{p8}{}_{AB} (A_{pmn} - 2B_{p[m} A_{n]}) \\ &\quad + V^{78}{}_{AB} B_{mn} + 2\tilde{V}_{[m|7}{}_{AB} A_{|n]} \end{aligned}$$

# “dressing up” process

[ vielbein and scalars ]

$$ds_4^2(x, y) = ds_4^2(x)$$

$$V^{m8 AB}(x, y) = \frac{1}{2} g K_{IJ}^m(y) \eta_i^A(y) \eta_j^B(y) \mathcal{V}^{IJ ij}(x) ,$$

$$V^{78 AB}(x, y) = -\mu_I(y) \eta_i^A(y) \eta_j^B(y) \mathcal{V}^{I8 ij}(x) ,$$

$$\tilde{V}_{mn}^{AB}(x, y) = \frac{1}{4} K_{mn}^{IJ}(y) \eta_i^A(y) \eta_j^B(y) \tilde{\mathcal{V}}_{IJ}^{ij}(x) ,$$

$$\tilde{V}_{m7}^{AB}(x, y) = -g^{-1} (\partial_m \mu^I)(y) \eta_i^A(y) \eta_j^B(y) \tilde{\mathcal{V}}_{I8}^{ij}(x) ,$$

[ two-forms ]

$$C_{\mu\nu m}{}^8(x, y) = -g^{-1} (\mu_I \partial_m \mu^J)(y) \mathcal{B}_{\mu\nu J}{}^I(x)$$

$$C_{\mu\nu 7}{}^8(x, y) = \mu_I(y) \mathcal{B}_{\mu\nu}{}^I(x)$$

[ three-form ]

$$C_{\mu\nu\rho}{}^{88}(x, y) = (\mu_I \mu_J)(y) \mathcal{C}_{\mu\nu\rho}{}^{IJ}(x)$$

[ vectors ]

$$C_\mu{}^{m8}(x, y) = \frac{1}{2} g K_{IJ}^m(y) \mathcal{A}_\mu{}^{IJ}(x) \quad , \quad C_\mu{}^{78}(x, y) = -\mu_I(y) \mathcal{A}_\mu{}^I(x) ,$$

$$\tilde{C}_{\mu mn}(x, y) = \frac{1}{4} K_{mn}^{IJ}(y) \tilde{\mathcal{A}}_{\mu IJ}(x) \quad , \quad \tilde{C}_{\mu m7}(x, y) = -g^{-1} (\partial_m \mu^I)(y) \tilde{\mathcal{A}}_{\mu I}(x)$$

◆  $S^6$  geometrical data : embedding coordinates  $\mu^I$ , Killing vectors  $K_{IJ}^m$  and tensors  $K_{IJ}^{mn}$

# 10D SUSY transformations

$$\delta C_\mu^{I8} = i V^{I8}{}_{AB} \left( \bar{\epsilon}^A \psi_\mu^B + \frac{1}{2\sqrt{2}} \bar{\epsilon}_C \gamma_\mu \chi^{ABC} \right) + \text{h.c.} ,$$

$$\delta \tilde{C}_{\mu IJ} = -i V_{IJ}{}_{AB} \left( \bar{\epsilon}^A \psi_\mu^B + \frac{1}{2\sqrt{2}} \bar{\epsilon}_C \gamma_\mu \chi^{ABC} \right) + \text{h.c.} ,$$

$$\begin{aligned} \delta C_{\mu\nu I}{}^8 &= \left[ \frac{2}{3} (V^{J8}{}_{BC} \tilde{V}_{IJ}{}^{AC} + \tilde{V}_{IJ}{}_{BC} V^{J8AC}) \bar{\epsilon}_A \gamma_{[\mu} \psi_{\nu]}^B \right. \\ &\quad \left. + \frac{\sqrt{2}}{3} V^{J8}{}_{AB} \tilde{V}_{IJCD} \bar{\epsilon}^{[A} \gamma_{\mu\nu} \chi^{BCD]} + \text{h.c.} \right] - C_{[\mu}{}^{J8} \delta \tilde{C}_{\nu]IJ} - \tilde{C}_{[\mu|IJ} \delta C_{|\nu]}{}^{J8} , \end{aligned}$$

$$\begin{aligned} \delta C_{\mu\nu\rho}{}^{88} &= \left[ \frac{4i}{7} V^{I8}{}_{BD} (V^{J8DC} \tilde{V}_{IJAC} + \tilde{V}_{IJ}{}^{DC} V^{J8AC}) \bar{\epsilon}^A \gamma_{[\mu\nu} \psi_{\rho]}^B \right. \\ &\quad \left. - i \frac{\sqrt{2}}{3} V^{I8AE} V^{J8}{}_{[EB|} \tilde{V}_{IJ|CD]} \bar{\epsilon}_A \gamma_{\mu\nu\rho} \chi^{BCD} + \text{h.c.} \right] \\ &\quad + 3 C_{[\mu\nu|I}{}^8 \delta C_{|\rho]}{}^{I8} - C_{[\mu}{}^{I8} (C_{\nu}{}^{J8} \delta \tilde{C}_{\rho]IJ} + \tilde{C}_{\nu|IJ} \delta C_{|\rho]}{}^{J8}) . \end{aligned}$$



# Freund-Rubin term

[ Freund & Rubin '80 ]

- By looking at the RR field strength  $\hat{F}_{(4)} = \mathcal{H}_{(4)}^{IJ} \mu_I \mu_J + \dots$ , one immediately identifies the Freund-Rubin term

$$\begin{aligned} \mathcal{H}_{(4)}^{IJ} \mu_I \mu_J = & -\frac{1}{3} g^{-1} V \text{vol}_4 + \frac{1}{84} g^{-1} \left( D\mathcal{H}_{(3)} - 7 \mathcal{H}_{(2)}^{IJ} \wedge \tilde{\mathcal{H}}_{(2)IJ} - 7 \mathcal{H}_{(2)}^I \wedge \tilde{\mathcal{H}}_{(2)I} \right) \\ & - \frac{1}{2} g^{-1} \left( D\mathcal{H}_{(3)I}{}^J - \mathcal{H}_{(2)}^{JK} \wedge \tilde{\mathcal{H}}_{(2)IK} - \mathcal{H}_{(2)}^J \wedge \tilde{\mathcal{H}}_{(2)I} \right) \mu^I \mu_J , \end{aligned}$$

**NOTE:** We have expressed the EOMs for the scalars as BI for the three-form field strengths of the tensor hierarchy.

- At a critical point of  $V$  one has  $\hat{F}_{(4)} = -\frac{1}{3g} V \text{vol}_4 + \dots$ , and the  $S^6$  dependence drops out

[ see also Godazgar, Godazgar, Krueger & Nicolai '15 ]