

# ABJM Baryon Stability at Finite 't Hooft Coupling

Yolanda Lozano (U. Oviedo)

Santiago de Compostela, October 2011

- **Motivation:** Study the stability of non-singlet baryon vertex-like configurations in ABJM
- **How:** Existence of a classical solution + stability analysis
  - Probe brane approx: Valid at **strong 't Hooft coupling**
  - Dissolve D0's  $\rightarrow$  Microscopical description: Valid at **finite 't Hooft coupling**
- **Results:**
  - Non-singlet stable baryons at **finite 't Hooft coupling**
  - Flat  $B_2$  required by Freed-Witten anomaly
  - New higher curvature dielectric couplings

(Based on arXiv:1105.0939 [hep-th], JHEP, with M. Picos, K. Sfetsos, K. Siampas)

# I. Introduction

$AdS_4/CFT_3$  relates the **Type IIA superstring** on  $AdS_4 \times CP^3$  to the  $\mathcal{N} = 6$  **Chern-Simons matter theory** with gauge group  $U(N)_k \times U(N)_{-k}$  known as **ABJM**.

- Good description when  $N^{1/5} \ll k$ .
- Like  $AdS_5/CFT_4$  it is a **strong weak coupling duality**, with ‘t Hooft coupling  $\lambda = N/k$ :
  - The string background describes the ‘t Hooft limit of the theory:  $N, k \rightarrow \infty$  with  $\lambda = N/k$  fixed
  - IIA weakly curved when  $k \ll N$  (large ‘t Hooft coupling)
- 3 dimensions  $\rightarrow$  Applications in condensed matter

## 2. Particle-like branes in ABJM

$CP^3$  has  $H^q(CP^3) = \mathbb{Z}$  for even  $q \Rightarrow$  D2, D4 and D6

particle-like branes wrapping topologically non-trivial cycles

## 2. Particle-like branes in ABJM

$CP^3$  has  $H^q(CP^3) = \mathbb{Z}$  for even  $q \Rightarrow$  D2, D4 and D6

particle-like branes wrapping topologically non-trivial cycles



### Interpretation in the dual CFT:

- D6 wrapped on the  $CP^3$ : Analogous to the baryon vertex in  $AdS_5 \times S^5$ .  $F_6$  flux  $\Rightarrow$  Tadpole that has to be cancelled with N F-strings ending on it  $\leftrightarrow$  N external quarks on the boundary of  $AdS_4$

# The baryon vertex in $AdS_5 \times S^5$

Gauge invariant coupling of N external quarks

Through AdS/CFT external quarks are regarded as endpoints of F-strings in AdS



Baryon vertex in the gravity side: D5-brane wrapped on the 5-sphere (Witten'98):

$$S_{CS} = 2\pi T_5 \int_{\mathbb{R} \times S^5} P[F_5] \wedge A = N T_{F1} \int_{\mathbb{R}} dt A_t$$

N charge cancelled by N F-strings ending on the 5-brane

Dual configuration on the CFT side:  $N$  Wilson lines ending on an epsilon tensor  $\longleftrightarrow$  Bound state of  $N$  quarks.

However, within the gauge/gravity correspondence it is possible to construct bound states of  $l$  quarks with  $l < N$  (non-singlets) (Brandhuber, Itzhaki, Sonnenschein, Yankielowicz'98; Imamura'98)

Dual configuration on the CFT side:  $N$  Wilson lines ending on an epsilon tensor  $\longleftrightarrow$  Bound state of  $N$  quarks.

However, within the gauge/gravity correspondence it is possible to construct bound states of  $l$  quarks with  $l < N$  (non-singlets) (Brandhuber, Itzhaki, Sonnenschein, Yankielowicz'98; Imamura'98)

In  $AdS_4 \times CP^3$ : A D6-brane wrapped on the  $CP^3$ :

$$S_{CS} = 2\pi T_6 \int_{\mathbb{R} \times CP^3} P[F_6] \wedge A = N T_{F1} \int dt A_t$$

Cancel this charge with the charge induced by the endpoints of  $N$  open F-strings stretching between the D6 and the boundary of AdS



Dual configuration on the CFT side:  $N$  Wilson lines ending on an epsilon tensor  $\longleftrightarrow$  Bound state of  $N$  quarks.

However, within the gauge/gravity correspondence it is possible to construct bound states of  $l$  quarks with  $l < N$  (non-singlets) (Brandhuber, Itzhaki, Sonnenschein, Yankielowicz'98; Imamura'98)

In  $AdS_4 \times CP^3$ : A D6-brane wrapped on the  $CP^3$ :

$$S_{CS} = 2\pi T_6 \int_{\mathbb{R} \times CP^3} P[F_6] \wedge A = N T_{F1} \int dt A_t$$

Cancel this charge with the charge induced by the endpoints of  $N$  open F-strings stretching between the D6 and the boundary of AdS

Non-singlets?

## 2. Particle-like branes in ABJM

$CP^3$  has  $H^q(CP^3) = \mathbb{Z}$  for even  $q \Rightarrow$  D2, D4 and D6

particle-like branes wrapping topologically non-trivial cycles

### Interpretation in the dual CFT:

- D6 wrapped on the  $CP^3$ : Analogous to the baryon vertex in  $AdS_5 \times S^5$ .  $F_6$  flux  $\Rightarrow$  Tadpole that has to be cancelled with N F-strings ending on it  $\leftrightarrow$  N external quarks on the boundary of  $AdS_4$
- D2 wrapped on a  $CP^1 \subset CP^3$ : 't Hooft monopole.  
 $F_2$  flux  $\Rightarrow$  Tadpole that has to be cancelled with  $k$  F-strings

But  $k$  Wilson lines cannot end on an epsilon tensor

If one forms the symmetric product only the endpoint of the Wilson lines is observable and the product behaves like a 't Hooft operator creating one unit of magnetic flux at a point (ABJM)  $\rightarrow$  't Hooft monopole

If one forms the symmetric product only the endpoint of the Wilson lines is observable and the product behaves like a 't Hooft operator creating one unit of magnetic flux at a point (ABJM)  $\rightarrow$  't Hooft monopole

- **D4 wrapped on a  $CP^2 \subset CP^3$ : Di-baryon**

It does not capture the background fluxes.

Same baryon charge and dimension than di-baryon:

Baryon charge  $N$ ,  $m_{D4}L = N \Rightarrow \Delta = \frac{m_{D4}L}{2} = \frac{N}{2}$

$\Rightarrow$  Dual configuration composed of  $N$  chirals

Di-baryon operator:  $O^{D4} = \epsilon_{i_1 \dots i_N} \epsilon^{j_1 \dots j_N} A_{j_1}^{i_1} \dots A_{j_N}^{i_N}$

If one forms the symmetric product only the endpoint of the Wilson lines is observable and the product behaves like a 't Hooft operator creating one unit of magnetic flux at a point (ABJM)  $\rightarrow$  't Hooft monopole

- D4 wrapped on a  $CP^2 \subset CP^3$ : Di-baryon

It does not capture the background fluxes.

Same baryon charge and dimension than di-baryon:

Baryon charge N,  $m_{D4}L = N \Rightarrow \Delta = \frac{m_{D4}L}{2} = \frac{N}{2}$

$\Rightarrow$  Dual configuration composed of N chirals

Di-baryon operator:  $O^{D4} = \epsilon_{i_1 \dots i_N} \epsilon^{j_1 \dots j_N} A_{j_1}^{i_1} \dots A_{j_N}^{i_N}$

These configurations admit a natural generalization by allowing non-trivial worldvolume gauge fluxes:

### 3. Add a magnetic flux

(Gutiérrez, Y.L., Rodríguez-Gómez'10)

Candidates for **holographic anyons in ABJM** (Kawamoto, Lin'09)  
(anyonic phase associated to the FI attached to the baryons surrounding the D0's dissolved)

### 3. Add a magnetic flux

(Gutiérrez, Y.L., Rodríguez-Gómez'10)

Candidates for **holographic anyons in ABJM** (Kawamoto, Lin'09)  
(anyonic phase associated to the FI attached to the baryons surrounding the D0's dissolved)

A **non-trivial flux** adds lower dim brane charges and **modifies the way the branes capture the background fluxes.**

### 3. Add a magnetic flux

(Gutiérrez, Y.L., Rodríguez-Gómez'10)

Candidates for **holographic anyons in ABJM** (Kawamoto, Lin'09)  
(anyonic phase associated to the FI attached to the baryons surrounding the D0's dissolved)

A **non-trivial flux** adds lower dim brane charges and **modifies the way the branes capture the background fluxes**.

For example, for  $F = \mathcal{N}J$  the D4 captures the  $F_2$  flux and develops a **tadpole**  $\Rightarrow$  **F-strings** ending on it :

$$S_{CS} = \frac{1}{2}(2\pi)^2 T_4 \int_{\mathbb{R} \times CP^2} P[F_2] \wedge F \wedge A = \frac{k\mathcal{N}}{2} T_{F1} \int dt A_t$$



### 3. Add a magnetic flux

(Gutiérrez, Y.L., Rodríguez-Gómez'10)

Candidates for **holographic anyons in ABJM** (Kawamoto, Lin'09)  
(anyonic phase associated to the FI attached to the baryons surrounding the D0's dissolved)

A **non-trivial flux** adds lower dim brane charges and **modifies the way the branes capture the background fluxes**.

For example, for  $F = \mathcal{N}J$  the D4 captures the  $F_2$  flux and develops a **tadpole**  $\Rightarrow$  **F-strings** ending on it :

$$S_{CS} = \frac{1}{2}(2\pi)^2 T_4 \int_{\mathbb{R} \times CP^2} P[F_2] \wedge F \wedge A = \frac{k\mathcal{N}}{2} T_{F1} \int dt A_t$$

$\rightarrow$  **Baryon vertex-like configurations**

### 3. Add a magnetic flux

(Gutiérrez, Y.L., Rodríguez-Gómez'10)

Candidates for **holographic anyons in ABJM** (Kawamoto, Lin'09)  
(anyonic phase associated to the FI attached to the baryons surrounding the D0's dissolved)

A **non-trivial flux** adds lower dim brane charges and **modifies the way the branes capture the background fluxes**.

For example, for  $F = \mathcal{N}J$  the D4 captures the  $F_2$  flux and develops a **tadpole**  $\Rightarrow$  **F-strings** ending on it :

$$S_{CS} = \frac{1}{2}(2\pi)^2 T_4 \int_{\mathbb{R} \times CP^2} P[F_2] \wedge F \wedge A = \frac{k\mathcal{N}}{2} T_{F1} \int dt A_t$$

$\rightarrow$  **Baryon vertex-like configurations**

**The magnetic flux modifies the dynamics as well:**

## 4. Gauge/gravity calculation of the energy

(Brandhuber, Itzhaki, Sonnenschein, Yankielowicz'98; Imamura'98; Maldacena'98)

Consider a uniform distribution of strings on a  $CP^{\frac{p}{2}}$  shell  
with  $p = 2, 4, 6$

Non-SUSY but we can ignore the backreaction

## 4. Gauge/gravity calculation of the energy

(Brandhuber, Itzhaki, Sonnenschein, Yankielowicz'98; Imamura'98; Maldacena'98)

Consider a uniform distribution of strings on a  $CP^{\frac{p}{2}}$  shell  
with  $p = 2, 4, 6$

Non-SUSY but we can ignore the backreaction


In the probe brane approx, with  $F = \mathcal{N}J$ ,  $S = S_{Dp} + S_{qF1}$ :

$$S_{Dp} = -Q_p \int dt \frac{2\rho}{L}, \quad Q_p = \frac{T_p}{g_s} \text{Vol}(CP^{\frac{p}{2}}) (L^4 + (2\pi\mathcal{N})^2)^{\frac{p}{4}}$$

$$S_{qT_{F1}} = -q T_{F1} \int dt dr \sqrt{\frac{16\rho^4}{L^4} + \rho'^2},$$

where we have taken  $\tau = t$  and  $\rho = \rho(r)$   
 $\sigma = r$

$\Leftrightarrow$  Radially symmetric distribution on a circle of radius  $l$



$$\rho(0) = \rho_0$$

$$\rho(l) = \infty$$

• ... Quarks

## 4. Gauge/gravity calculation of the energy

Consider a uniform distribution of strings on a  $CP^{\frac{p}{2}}$  shell  
with  $p = 2, 4, 6$

Non-SUSY but we can ignore the backreaction

In the probe brane approx, with  $F = \mathcal{N}J$ ,  $S = S_{Dp} + S_{qF1}$ :

$$S_{Dp} = -Q_p \int dt \frac{2\rho}{L}, \quad Q_p = \frac{T_p}{g_s} \text{Vol}(CP^{\frac{p}{2}}) (L^4 + (2\pi\mathcal{N})^2)^{\frac{p}{4}}$$

$$S_{qT_{F1}} = -q T_{F1} \int dt dr \sqrt{\frac{16\rho^4}{L^4} + \rho'^2}$$

Bulk equation of motion: 
$$\frac{\rho^4}{\sqrt{\frac{16\rho^4}{L^4} + \rho'^2}} = c$$

Boundary equation of motion: 
$$\frac{\rho'_0}{\sqrt{\frac{16\rho_0^4}{L^4} + \rho_0'^2}} = \frac{2Q_p}{L q T_{F1}}$$

Define  $\sqrt{1 - \beta^2} = \frac{2Q_p}{L_q T_{F1}}$  with  $\beta \in [0, 1]$

The two equations can be combined into:

$$\frac{\rho^4}{\sqrt{\frac{16\rho^4}{L^4} + \rho'^2}} = \frac{1}{4} \beta \rho_0^2 L^2$$

Integrating:      Size of the configuration:

$$\ell = \frac{L^2}{4\rho_0} \int_1^\infty dz \frac{\beta}{z^2 \sqrt{z^4 - \beta^2}}$$

Same form for the baryon vertex in  $AdS_5 \times S^5$

Same dependence on  $L^2$  in  $AdS_5 \times S^5$  : Prediction of AdS/CFT  
for the strong coupling behavior of the CS theory

## On-shell energy:

$$E = E_{Dp} + E_{qF1} = qT_{F1}\rho_0 \left( \sqrt{1 - \beta^2} + \int_1^\infty dz \frac{z^2}{\sqrt{z^4 - \beta^2}} \right)$$

## Binding energy:

$$E_{\text{bin}} = qT_{F1}\rho_0 \left( \sqrt{1 - \beta^2} + \int_1^\infty dz \left[ \frac{z^2}{\sqrt{z^4 - \beta^2}} - 1 \right] - 1 \right)$$

where we have subtracted the energy of the constituents  
(when the brane is located in  $\rho_0 = 0$  the strings become  
radial and correspond to free quarks)

- $E_{\text{bin}}$  negative and decreases monotonically with  $\beta$
- $E_{\text{bin}} = 0$  for  $\beta = 0$  (q free radial strings stretching from  $\rho_0$  to  $\infty$  plus a Dp-brane at  $\rho_0$ ) (only for non-zero magnetic flux)



As a function of  $\ell$  :

$$E_{\text{bin}} = -f(\beta) \frac{(g_s N)^{2/5}}{\ell} \quad \text{with} \quad f(\beta) \geq 0$$

$\Rightarrow$  - The configuration is stable

-  $E_{\text{bin}} \sim 1/\ell$  dictated by conformal invariance

- As a function of the 't Hooft coupling,  $\lambda = N/k$ ,

$E_{\text{bin}} \sim \sqrt{\lambda}$ , as in  $AdS_5 \Rightarrow$  **Non-trivial prediction for the non-perturbative regime of the CS theory**  
(Mariño, Putrov'09)

In fact, since the D4 wraps a non-spin manifold it must carry  $F_{FW} = J$  due to the **Freed-Witten anomaly** (Freed, Witten'99)

$\Rightarrow$  A **flat half-integer**  $B_2$  has to be switched on, such that

$$\mathcal{F} = F_{FW} + \frac{1}{2\pi} B_2 = 0$$

Then  $Q_p = \frac{T_p}{g_s} \text{Vol}(CP^{\frac{p}{2}}) (L^4 + (2\pi)^2 (\mathcal{N} - 1)^2)^{\frac{p}{4}}$  for  $p = 2, 6$




For the D6 there are **new CS terms**:

$$\int_{\mathbb{R} \times CP^3} P[F_2] \wedge B_2 \wedge B_2 \wedge A, \quad \int_{\mathbb{R} \times CP^3} P[F_2] \wedge F \wedge B_2 \wedge A$$

that **modify the number of F-strings**. First is cancelled with

$$S_{h.c.}^{D6} = \frac{3}{2} (2\pi)^5 T_6 \int C_1 \wedge F \wedge \sqrt{\frac{\hat{A}(T)}{\hat{A}(N)}} \quad (\text{Aharony, Hashimoto, Hirano, Ouyang'09})$$

$$\frac{2Q_p}{L q T_{F1}} \leq 1 \Rightarrow \text{Bound on the magnetic flux}$$



Binding energy per string

$q$  : Number of strings

D-brane	$q$
D2	$k$
D4	$k \frac{\mathcal{N}}{2}$
D6	$N + k \frac{\mathcal{N}(\mathcal{N} - 2)}{8}$


## 5. Reduce the number of fundamental strings

In  $AdS_5 \times S^5$  : Baryon vertex classical solutions with number of quarks  $5N/8 \leq l \leq N$  (**non-singlet**)

(Brandhuber, Itzhaki, Sonnenschein, Yankielowicz'98;  
Imamura'98)

Stable against fluctuations for  $0.813N \leq l \leq N$   
(Sfetsos, Siampos'08)

In  $AdS_4 \times CP^3$  :



## Analysis of the equations of motion:

→ Classical solutions exist for  $\frac{q}{2}(1 + \sqrt{1 - \beta^2}) \leq l \leq q$

→ Bound on the magnetic flux

$(l, \mathcal{N})$  parameter space bounded by the values for which the baryon vertex reduces to free quarks

## Analysis of the equations of motion:

→ Classical solutions exist for  $\frac{q}{2}(1 + \sqrt{1 - \beta^2}) \leq l \leq q$

→ Bound on the magnetic flux

$(l, \mathcal{N})$  parameter space bounded by the values for which the baryon vertex reduces to free quarks

## Stability analysis

Instabilities emerge from longitudinal fluctuations of the strings  
(Sfetsos, Siampos'08)

Perturb the classical solution:  $\delta x^\mu(t, \rho) = \delta x^\mu(\rho)e^{-i\omega t}$  for  $x^\mu = r, \theta$ , expand the Nambu-Goto action to quadratic order and study the zero mode problem

Bound for the number of F-strings coming from stability:

$$l \geq \frac{q}{1 + \gamma_c} (1 + \sqrt{1 - \beta^2}) \quad \gamma_c = 0.538$$

Bound for the number of F-strings coming from stability:

$$l \geq \frac{q}{1 + \gamma_c} (1 + \sqrt{1 - \beta^2}) \quad \gamma_c = 0.538$$

More restrictive than the bound imposed by the existence of a classical solution:

$$l \geq \frac{q}{2} (1 + \sqrt{1 - \beta^2})$$



Bound for the number of F-strings coming from stability:

$$l \geq \frac{q}{1 + \gamma_c} (1 + \sqrt{1 - \beta^2}) \quad \gamma_c = 0.538$$

More restrictive than the bound imposed by the existence of a classical solution:

$$l \geq \frac{q}{2} (1 + \sqrt{1 - \beta^2})$$

$\Rightarrow$  Non-singlet states exist for  $\lambda \gg 1$

Bound for the number of F-strings coming from stability:

$$l \geq \frac{q}{1 + \gamma_c} (1 + \sqrt{1 - \beta^2}) \quad \gamma_c = 0.538$$

More restrictive than the bound imposed by the existence of a classical solution:

$$l \geq \frac{q}{2} (1 + \sqrt{1 - \beta^2})$$

$\Rightarrow$  Non-singlet states exist for  $\lambda \gg 1$

Can we reach the finite 't Hooft coupling region?

Bound for the number of F-strings coming from stability:

$$l \geq \frac{q}{1 + \gamma_c} (1 + \sqrt{1 - \beta^2}) \quad \gamma_c = 0.538$$

More restrictive than the bound imposed by the existence of a classical solution:

$$l \geq \frac{q}{2} (1 + \sqrt{1 - \beta^2})$$

$\Rightarrow$  Non-singlet states exist for  $\lambda \gg 1$

Can we reach the finite 't Hooft coupling region?

Microscopical description in terms of fuzzy  $CP^{\frac{p}{2}}$  built up out of  $n$  dielectric D0-branes valid when  $N \ll n^{\frac{4}{p}} k$

## 6. The microscopical description

A non-trivial magnetic flux induces D0-brane charge on the Dp

⇒ Complementary description in terms of multiple D0  
expanded into fuzzy  $CP^{\frac{p}{2}}$  by Myers dielectric effect

## 6. The microscopical description

A non-trivial magnetic flux induces D0-brane charge on the Dp

⇒ Complementary description in terms of multiple D0  
expanded into fuzzy  $CP^{\frac{p}{2}}$  by Myers dielectric effect

- Macroscopical description valid in the sugra limit:

$$L \gg 1 \Leftrightarrow N \gg k$$

## 6. The microscopical description

A non-trivial magnetic flux induces D0-brane charge on the Dp

⇒ Complementary description in terms of multiple D0  
expanded into fuzzy  $CP^{\frac{p}{2}}$  by Myers dielectric effect

- Macroscopical description valid in the sugra limit:

$$L \gg 1 \Leftrightarrow N \gg k$$

- Micro when

$$\frac{\text{Vol}(CP^{\frac{p}{2}})}{n} \ll l_s^p \Leftrightarrow N \ll n^{\frac{4}{p}} k$$

## 6. The microscopical description

A non-trivial magnetic flux induces D0-brane charge on the Dp

⇒ Complementary description in terms of multiple D0  
expanded into fuzzy  $CP^{\frac{p}{2}}$  by Myers dielectric effect

- Macroscopical description valid in the sugra limit:

$$L \gg 1 \Leftrightarrow N \gg k$$

- Micro when

$$\frac{\text{Vol}(CP^{\frac{p}{2}})}{n} \ll l_s^p \Leftrightarrow N \ll n^{\frac{4}{p}} k$$

⇒ It allows to explore the region of finite  $\lambda$

## 6. The microscopical description

A non-trivial magnetic flux induces D0-brane charge on the Dp

⇒ Complementary description in terms of multiple D0  
expanded into fuzzy  $CP^{\frac{p}{2}}$  by Myers dielectric effect

- Macroscopical description valid in the sugra limit:

$$L \gg 1 \Leftrightarrow N \gg k$$

- Micro when

$$\frac{\text{Vol}(CP^{\frac{p}{2}})}{n} \ll l_s^p \Leftrightarrow N \ll n^{\frac{4}{p}} k$$

⇒ It allows to explore the region of finite  $\lambda$

Complementary for finite  $n$

Should agree in the large  $n$  limit



## 6.1. Fuzzy CP

- $CP^{\frac{p}{2}}$ : coset manifold  $\frac{SU(\frac{p}{2} + 1)}{U(\frac{p}{2})}$ . Submanifold of  $\mathbb{R}^{\frac{p^2}{4} + p}$  determined by the constraints:

$$\sum_{i=1}^{\frac{p^2}{4} + p} (x^i)^2 = 1, \quad \sum_{j,k=1}^{\frac{p^2}{4} + p} d^{ijk} x^j x^k = \frac{\frac{p}{2} - 1}{\sqrt{\frac{p}{4}(\frac{p}{2} + 1)}} x^i$$

→  $p$  dimensional manifold

Fubini-Study metric of the  $CP^{\frac{p}{2}}$  given by

$$ds_{CP^{\frac{p}{2}}}^2 = \frac{p}{4(\frac{p}{2} + 1)} \sum_{i=1}^{\frac{p^2}{4} + p} (dx^i)^2$$

## 6.1. Fuzzy CP

- $CP^{\frac{p}{2}}$ : coset manifold  $\frac{SU(\frac{p}{2} + 1)}{U(\frac{p}{2})}$ . Submanifold of  $\mathbb{R}^{\frac{p^2}{4} + p}$  determined by the constraints:

$$\sum_{i=1}^{\frac{p^2}{4} + p} (x^i)^2 = 1, \quad \sum_{j,k=1}^{\frac{p^2}{4} + p} d^{ijk} x^j x^k = \frac{\frac{p}{2} - 1}{\sqrt{\frac{p}{4}(\frac{p}{2} + 1)}} x^i$$

→  $p$  dimensional manifold

Fubini-Study metric of the  $CP^{\frac{p}{2}}$  given by

$$ds_{CP^{\frac{p}{2}}}^2 = \frac{p}{4(\frac{p}{2} + 1)} \sum_{i=1}^{\frac{p^2}{4} + p} (dx^i)^2$$

- Matrix level definition  $\leftrightarrow$  Fuzzy  $CP^{\frac{p}{2}}$  :

$$X^i = \frac{1}{\sqrt{C_n}} T^i, \quad T^i : \text{generators of } SU(\frac{p}{2} + 1) \text{ in the } (m, 0) \text{ irrep}$$

Substituting in **Myers action for D0-branes**:

$$S_{DBI} = - \int \text{STr} \left\{ e^{-\phi} \sqrt{|\det \left( P[E_{\mu\nu} + E_{\mu i} (Q^{-1} - \delta)^i_j E^{jk} E_{k\nu}] \right) \det Q|} \right\}$$

$$Q^i_j = \delta^i_j + \frac{i}{2\pi} [X^i, X^k] E_{kj}$$

## Substituting in Myers action for D0-branes:

$$S_{DBI} = - \int \text{STr} \left\{ e^{-\phi} \sqrt{|\det \left( P[E_{\mu\nu} + E_{\mu i} (Q^{-1} - \delta)^i_j E^{jk} E_{k\nu}] \right) \det Q|} \right\}$$

$$Q^i_j = \delta^i_j + \frac{i}{2\pi} [X^i, X^k] E_{kj}$$

$$\Rightarrow S_{nD0}^{DBI} = -\frac{n}{g_s} \left( 1 + \frac{L^4}{16\pi^2 m(m + \frac{p}{2} + 1)} \right)^{\frac{p}{4}} \int dt \frac{2\rho}{L}$$

## Substituting in Myers action for D0-branes:

$$S_{DBI} = - \int \text{STr} \left\{ e^{-\phi} \sqrt{|\det \left( P[E_{\mu\nu} + E_{\mu i} (Q^{-1} - \delta)^i_j E^{jk} E_{k\nu}] \right) \det Q|} \right\}$$

$$Q^i_j = \delta^i_j + \frac{i}{2\pi} [X^i, X^k] E_{kj}$$

$$\Rightarrow S_{nD0}^{DBI} = -\frac{n}{g_s} \left( 1 + \frac{L^4}{16\pi^2 m(m + \frac{p}{2} + 1)} \right)^{\frac{p}{4}} \int dt \frac{2\rho}{L}$$

$$n = \dim(m, 0), \quad n = \frac{\mathcal{N}^{\frac{p}{2}}}{2^{\frac{p}{2}} (\frac{p}{2})!} + \dots \Rightarrow m \sim \frac{\mathcal{N}}{2} \quad \text{for large } n$$

and  $S_{nD0}^{DBI}$  exactly reproduces the macroscopical result:

$$S_{Dp} = -Q_p \int dt \frac{2\rho}{L}, \quad Q_p = \frac{T_p}{g_s} \text{Vol}(CP^{\frac{p}{2}}) (L^4 + (2\pi\mathcal{N})^2)^{\frac{p}{4}}$$

## 6.2. The F-strings from the Dp to the boundary of AdS

CS action for coincident branes:

$$S_{CS} = \int \text{STr} \left\{ P \left( e^{\frac{i}{2\pi} (i_X i_X)} \sum_q C_q e^{B_2} \right) e^{2\pi F} \right\}$$

Dependence of the background potentials on the non-Abelian scalars:

$$C_q(t, X) = C_q(t) + X^k \partial_k C_q(t) + \frac{1}{2} X^l X^k \partial_l \partial_k C_q(t) + \dots$$

## 6.2. The F-strings from the Dp to the boundary of AdS

CS action for coincident branes:

$$S_{CS} = \int \text{STr} \left\{ P \left( e^{\frac{i}{2\pi} (i_X i_X)} \sum_q C_q e^{B_2} \right) e^{2\pi F} \right\}$$

Dependence of the background potentials on the non-Abelian scalars:

$$C_q(t, X) = C_q(t) + X^k \partial_k C_q(t) + \frac{1}{2} X^l X^k \partial_l \partial_k C_q(t) + \dots$$

For example:

$$\begin{aligned} S_{CS_2} &= -\frac{i}{(2\pi)^2} \int \text{STr} \{ (i_X i_X)^3 F_6 \wedge A \} = \\ &= N \left( m(m+4) \right)^{-3/2} \frac{(m+3)!}{m!} \int dt A_t \end{aligned}$$

## 6.2. The F-strings from the Dp to the boundary of AdS

CS action for coincident branes:

$$S_{CS} = \int \text{STr} \left\{ P \left( e^{\frac{i}{2\pi} (i_X i_X)} \sum_q C_q e^{B_2} \right) e^{2\pi F} \right\}$$

Dependence of the background potentials on the non-Abelian scalars:

$$C_q(t, X) = C_q(t) + X^k \partial_k C_q(t) + \frac{1}{2} X^l X^k \partial_l \partial_k C_q(t) + \dots$$

For example:

$$\begin{aligned} S_{CS_2} &= -\frac{i}{(2\pi)^2} \int \text{STr} \{ (i_X i_X)^3 F_6 \wedge A \} = \\ &= N \left( m(m+4) \right)^{-3/2} \frac{(m+3)!}{m!} \int dt A_t \rightarrow N \int dt A_t \quad \text{for } m \gg 1 \end{aligned}$$



$S_{CS_1} = i \int \text{STr}\{(i_X i_X) F_2 \wedge A\}$  gives the FI charge prop. to  $k$

$S_{CS_1} = i \int \text{STr}\{(i_X i_X) F_2 \wedge A\}$  gives the FI charge prop. to  $k$

### 6.3. The flat half-integer NS-NS 2-form

Introduced macroscopically to cancel the flux of the Freed-Witten vector field, such that  $\mathcal{F} = F_{FW} + \frac{1}{2\pi} B_2 = 0$

Microscopically: We should find an obstacle to the expansion of the D0 into a fuzzy  $CP^2$  when  $B_2 = 0$ .

However,  $F_{FW}$  does not couple in the action for D0

How precisely  $B_2 \neq 0$  allows the construction of the  $CP^2$  ?

Macroscopically  $B_2$  modifies the BI action such that  $\mathcal{N} \rightarrow \mathcal{N} - 1$  (for the D2 and D6)

$\Rightarrow$  Microscopically we should include  $\frac{1}{m}$  corrections

To this order we find for  $B_2 = 0$ :

$$\mathcal{N} = 2m + \frac{p}{2} + 1 \Rightarrow \mathcal{N} \in 2\mathbb{Z} \quad \text{for } p=2,6$$

Whereas for  $B_2 \neq 0$ :

$$\mathcal{N}_{D2,D6} = 2m + \frac{p}{2} + 1 \quad \text{and} \quad \mathcal{N} \rightarrow \mathcal{N} - 1$$

$$\mathcal{N}_{D4} = 2m + \frac{p}{2} \Rightarrow \mathcal{N} \in 2\mathbb{Z} \quad \text{for all } p$$

Macroscopically  $B_2$  modifies the BI action such that  $\mathcal{N} \rightarrow \mathcal{N} - 1$  (for the D2 and D6)

$\Rightarrow$  Microscopically we should include  $\frac{1}{m}$  corrections

To this order we find for  $B_2 = 0$ :

$$\mathcal{N} = 2m + \frac{p}{2} + 1 \Rightarrow \mathcal{N} \in 2\mathbb{Z} \quad \text{for } p=2,6$$

Whereas for  $B_2 \neq 0$ :

$$\mathcal{N}_{D2,D6} = 2m + \frac{p}{2} + 1 \quad \text{and} \quad \mathcal{N} \rightarrow \mathcal{N} - 1$$

$$\mathcal{N}_{D4} = 2m + \frac{p}{2} \Rightarrow \mathcal{N} \in 2\mathbb{Z} \quad \text{for all } p$$

$\Rightarrow B_2 \neq 0$  to have  $\mathcal{N}$  properly quantized

Confirmed by the CS action (include as well

$$S_{CS_3} = -\frac{1}{2\pi} \int \text{STr} \left\{ (i_X i_X)^2 F_2 \wedge B_2 \wedge A \right\} )$$

Confirmed by the CS action (include as well

$$S_{CS_3} = -\frac{1}{2\pi} \int \text{STr} \left\{ (i_X i_X)^2 F_2 \wedge B_2 \wedge A \right\} )$$

Higher curvature couplings are needed in order to cancel the contribution of

$$S_{CS_4} = -\frac{i}{2} \frac{1}{(2\pi)^2} \int \text{STr} \left\{ (i_X i_X)^3 F_2 \wedge B_2 \wedge B_2 \wedge A \right\}$$

Confirmed by the CS action (include as well

$$S_{CS_3} = -\frac{1}{2\pi} \int \text{STr} \left\{ (i_X i_X)^2 F_2 \wedge B_2 \wedge A \right\} )$$

Higher curvature couplings are needed in order to cancel the contribution of

$$S_{CS_4} = -\frac{i}{2} \frac{1}{(2\pi)^2} \int \text{STr} \left\{ (i_X i_X)^3 F_2 \wedge B_2 \wedge B_2 \wedge A \right\}$$

$$\begin{aligned} S_{h.c.} &= -\frac{1}{2(2\pi)^2} \int_{\mathbb{R}} P \left[ (i_X i_X)^2 C_1 \wedge \sqrt{\frac{\hat{A}(T)}{\hat{A}(N)}} \right] = \\ &= -\frac{i}{(2\pi)^2} \int_{\mathbb{R}} \left[ (i_X i_X)^3 (F_2 \wedge \sqrt{\frac{\hat{A}(T)}{\hat{A}(N)}}) \right] A \end{aligned}$$

In general:

$$S_{h.c.} = T_p \int d^{p+1} \xi \text{STr} \left[ P \left( e^{\frac{i}{2\pi} (i_X i_X)} \sum_q C_q e^{B_2} \sqrt{\frac{\hat{A}(T)}{\hat{A}(N)}} \right) e^{2\pi F} \right]_{p+1}$$



In general:

$$S_{h.c.} = T_p \int d^{p+1} \xi \text{STr} \left[ P \left( e^{\frac{i}{2\pi} (i_X i_X)} \sum_q C_q e^{B_2} \sqrt{\frac{\hat{A}(T)}{\hat{A}(N)}} \right) e^{2\pi F} \right]_{p+1}$$

⇒ New dielectric couplings of RR-fields to derivatives of  $B_2$  and the metric through T-duality

(Becker, Guo, Robbins'10) (Garousi'10)

In general:

$$S_{h.c.} = T_p \int d^{p+1} \xi \text{STr} \left[ P \left( e^{\frac{i}{2\pi} (i_X i_X)} \sum_q C_q e^{B_2} \sqrt{\frac{\hat{A}(T)}{\hat{A}(N)}} \right) e^{2\pi F} \right]_{p+1}$$

⇒ New dielectric couplings of RR-fields to derivatives of  $B_2$  and the metric through T-duality

(Becker, Guo, Robbins'10) (Garousi'10)

Similar stability analysis

## 7. Conclusions

Stability of baryon vertex like configurations in  $AdS_4 \times CP^3$ :

- Condition for existence of a classical solution:

$$l \geq \frac{q}{2}(1 + \sqrt{1 - \beta^2})$$

- Stable when  $l \geq \frac{q}{1 + \gamma_c}(1 + \sqrt{1 - \beta^2})$   $\gamma_c = 0.538$

$\Rightarrow$  More restrictive

- Probe brane approx  $\Rightarrow$  Valid in the **SUGRA limit**  $\lambda \gg 1$
- For non-zero magnetic flux: D0-brane charge dissolved  $\Rightarrow$

Alternative description in terms of D0-branes expanded into fuzzy  $CP^{\frac{p}{2}}$  by Myers dielectric effect

## Microscopical description valid for finite 't Hooft coupling

- Expansion caused by a purely gravitational dielectric effect
- CS terms indicate the need to introduce F-strings
- Non-singlet classical stable solutions for finite  $\lambda$
- Prediction of new dielectric higher curvature couplings, with further implications through T and S duality

(Becker, Guo, Robbins'10) (Garousi'10)

## 8. Open questions

- New dielectric higher curvature couplings confirmed from string amplitudes?
- Can we extrapolate the micro results to  $\mathcal{N} \rightarrow 0$  ?
- What happens in theories with less susy, like  $AdS_5 \times T^{1,1}$  ?
- Include the backreaction  $\longleftrightarrow$  Look for supersymmetric spike solutions (partial studies in Kawamoto, Lin'09)

Marginal bound states?

Non-singlet states?

- Explore the finite temperature case

## 8. Open questions

- New dielectric higher curvature couplings confirmed from string amplitudes?
- Can we extrapolate the micro results to  $\mathcal{N} \rightarrow 0$  ?
- What happens in theories with less susy, like  $AdS_5 \times T^{1,1}$  ?
- Include the backreaction  $\longleftrightarrow$  Look for supersymmetric spike solutions (partial studies in Kawamoto, Lin'09)

Marginal bound states?

Non-singlet states?

- Explore the finite temperature case

Thanks!