

# Neutral Binuclear and Anionic Trinuclear Rhenium Carbonyl Complexes Containing Bridging Ligands Derived from 2-Amino- and 2-Mercaptopyridines

Javier A. Cabeza,<sup>\*,†</sup> Angela Llamazares, Víctor Riera, and Rajiv Trivedi

*Departamento de Química Orgánica e Inorgánica, Instituto de Química Organometálica "Enrique Moles", Universidad de Oviedo-CSIC, E-33071 Oviedo, Spain*

Fabrizia Grepioni<sup>\*,‡</sup>

*Dipartimento di Chimica "G. Ciamician", Università degli Studi di Bologna, via Selmi 2, I-40126 Bologna, Italy*

Received June 30, 1998

The reactions of the nitrile-stabilized complex  $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{11}(\text{MeCN})]$  (**1**) and of the unsaturated complex  $[\text{Re}_4(\mu\text{-H})_4(\text{CO})_{12}]$  (**2**) with 2-amino-6-methylpyridine (Hampy), 2-mercaptopyridine (Hmerpy), and 2-hydroxy-6-methylpyridine (Hhympy) have been studied. Complex **1** reacts slowly with Hampy and Hmerpy (refluxing toluene, 5–7 h) to give the binuclear derivatives  $[\text{Re}_2(\mu\text{-ampy})_2(\text{CO})_6]$  (**3**) and  $[\text{Re}_2(\mu\text{-merpy})_2(\text{CO})_6]$  (**4**), but no reaction is observed between **1** and Hhympy under analogous conditions. Complex **2** reacts with Hampy and Hmerpy (refluxing 1,2-dichloroethane, 10–20 min) to give the trinuclear anionic derivatives  $[\text{Re}_3(\mu\text{-H})_3(\mu_3\text{-ampy})(\text{CO})_9]^-$  (**5**) and  $[\text{Re}_3(\mu\text{-H})_3(\mu_3\text{-merpy})(\text{CO})_9]^-$  (**6**), which have been isolated as pyridinium ( $[\text{H}_2\text{ampy}]^+$ **5**) or triethylammonium ( $[\text{HNET}_3]^+$ **5**,  $[\text{HNET}_3]^+$ **6**) salts. Under analogous conditions, the reaction of **2** with Hhympy gives an inseparable mixture of compounds. The X-ray structures of  $[\text{HNET}_3]^+$ **5** and  $[\text{HNET}_3]^+$ **6** confirm that the three edges of the clusters are spanned by hydride ligands, that the ampy (in **5**) and merpy (in **6**) are attached to the three Re atoms, and that there are hydrogen-bonding interactions in the solid state between some carbonyl oxygen atoms and the NH fragments of the ampy ligand (in **5**) and the triethylammonium cation (in **5** and **6**).

## Introduction

To date, the carbonyl cluster compounds (three or more metal atoms connected by metal–metal bonds) known for rhenium are still very few<sup>1</sup> in comparison with those known for ruthenium and osmium.<sup>2</sup> This is mainly due to the lack of a convenient rhenium starting material. In fact, the derivative chemistry of the most readily available rhenium carbonyl compound,  $[\text{Re}_2\text{-}$

$(\text{CO})_{10}]$ , leads only to cluster derivatives of interest as starting materials in its reactions with hydrogen,<sup>3</sup> hydrides,<sup>4</sup> and hydroxide anions,<sup>5</sup> but the products are generally isolated in medium to low yields. For example,  $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{12}]$ , which is isoelectronic with the most important ruthenium and osmium starting clusters  $[\text{M}_3(\text{CO})_{12}]$  ( $\text{M} = \text{Ru}, \text{Os}$ ), can be prepared from the expensive  $[\text{Re}_2(\text{CO})_{10}]$  by either hydrogenation<sup>3a</sup> or by treatment with  $\text{Na}[\text{BH}_4]$  followed by acidification,<sup>4d,e</sup> but both procedures are rather tedious and the yields are only moderate (50% at best). Moreover,  $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{12}]$  is kinetically very stable, and its reactions require high temperatures (which promote changes in cluster and/or ligand integrity, often leading to mixtures

\* Corresponding author. Fax: +34-985103446. E-mail: jac@sauron.quimica.uniovi.es.

† E-mail: jac@sauron.quimica.uniovi.es.

‡ E-mail: grepioni@ciam.unibo.it.

(1) For comprehensive reviews on rhenium carbonyl cluster chemistry, see: (a) O'Connor, J. M. In *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1995; Vol. 6 (Casey, C. P., Ed.), p 173. (b) Henly, T. J. *Coord. Chem. Rev.* **1989**, *93*, 269.

(2) For comprehensive reviews on ruthenium and osmium carbonyl cluster chemistry, see: (a) Deeming, A. J. In *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1995; Vol. 7 (Shriver, D. F., Bruce, M. I., Eds.), p 684. (b) Smith, A. K. In *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1995; Vol. 7 (Shriver, D. F., Bruce, M. I., Eds.), p 804. (d) Pomeroy, R. K. In *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1995; Vol. 7 (Shriver, D. F., Bruce, M. I., Eds.), p 836. (e) Cifuentes, M. P.; Humphrey, M. G. In *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1995; Vol. 7 (Shriver, D. F., Bruce, M. I., Eds.), p 908.

(3) (a) Kaesz, H. D.; Knox, S. A. R.; Koepke, J. W.; Saillant, R. B. *J. Chem. Soc., Chem. Commun.* **1971**, 447. (b) Saillant, R. B.; Barceló, G.; Kaesz, H. D. *J. Am. Chem. Soc.* **1970**, *92*, 5739. (c) Johnson, R. J.; Kaesz, H. D. *Inorg. Synth.* **1978**, *18*, 60. (d) Wang, S. R.; Wang, S. L.; Cheng, C. P.; Yang, C. S. *J. Organomet. Chem.* **1992**, *431*, 215.

(4) (a) Huie, B. T.; Kirtley, S. W.; Knobler, C. B.; Kaesz, H. D. *J. Organomet. Chem.* **1981**, *213*, 45. (b) Bau, R.; Fontal, B.; Kaesz, H. D.; Churchill, M. R. *J. Am. Chem. Soc.* **1967**, *89*, 6374. (c) Kaesz, H. D.; Fontal, B.; Bau, R.; Kirtley, S. W.; Churchill, M. R. *J. Am. Chem. Soc.* **1969**, *91*, 1021. (d) Andrews, M. A.; Kirtley, S. W.; Kaesz, H. D. *Inorg. Synth.* **1977**, *17*, 66. (e) Huggins, D. K.; Fellmann, W.; Smith, J. M.; Kaesz, H. D. *J. Am. Chem. Soc.* **1964**, *86*, 4841.


(5) (a) Ciani, G.; Albano, V. G.; Immirzi, A. *J. Organomet. Chem.* **1976**, *121*, 237. (b) Bertolucci, A.; Freni, M.; Romiti, P.; Ciani, G.; Sironi, A.; Albano, V. G. *J. Organomet. Chem.* **1976**, *113*, C61. (c) Bertolucci, A.; Ciani, G.; Freni, M.; Romiti, P.; Albano, V. G.; Albinati, A. *J. Organomet. Chem.* **1976**, *117*, C37. (d) Freni, M.; Romiti, P.; D'Alfonso, G. *J. Organomet. Chem.* **1977**, *140*, 195.

of products)<sup>1</sup> or derivatization to the more reactive [Re<sub>3</sub>(μ-H)<sub>3</sub>(CO)<sub>10</sub>(MeCN)<sub>2</sub>].<sup>6</sup> In addition, although the derivative chemistry of the anionic unsaturated cluster [Re<sub>3</sub>(μ-H)<sub>4</sub>(CO)<sub>10</sub>]<sup>-</sup> has been extensively studied by Ciani, D'Alfonso, and co-workers,<sup>1</sup> its best preparation requires the hydrogenation at high pressure (100 atm) of [Re<sub>3</sub>(μ-H)<sub>2</sub>(CO)<sub>12</sub>]<sup>-</sup>,<sup>7</sup> which is made by deprotonation of [Re<sub>3</sub>(μ-H)<sub>3</sub>(CO)<sub>12</sub>].<sup>8</sup>

On the other hand, we (and other authors) have previously studied the reactivity of [Ru<sub>3</sub>(CO)<sub>12</sub>] with ortho-functionalized pyridines such as 2-amino-,<sup>9,10</sup> 2-mercapto-,<sup>10-12</sup> and 2-hydroxypyridines.<sup>9,10</sup> In the case of 2-aminopyridines, the products are trinuclear derivatives that contain the corresponding monodeprotonated ligand in a face-bridging mode, such as [Ru<sub>3</sub>(μ-H)(μ<sub>3</sub>-ampy)(CO)<sub>12</sub>] (Hampy = 2-amino-6-methylpyridine).<sup>9</sup> The presence of such bridging ligands has profound implications in the reactivity of the clusters, since they help maintain the cluster integrity and provide reaction pathways of low activation energy.<sup>13</sup> Over a hundred ruthenium carbonyl clusters containing μ<sub>3</sub>-2-aminopyridinate ligands have already been prepared,<sup>2a</sup> including the first cationic nonhydridic 48-electron triruthenium clusters,<sup>14</sup> some hexanuclear clusters,<sup>15</sup> and some efficient catalyst precursors for alkyne hydrogenation,<sup>16</sup> dimerization,<sup>17</sup> and polymerization<sup>17</sup> reactions.

We now describe some approaches to the synthesis of rhenium carbonyl cluster compounds containing bridging ligands derived from 2-amino-, 2-mercapto-, and 2-hydroxypyridines, using [Re<sub>3</sub>(μ-H)<sub>3</sub>(CO)<sub>11</sub>(MeCN)] (1) and [Re<sub>4</sub>(μ-H)<sub>4</sub>(CO)<sub>12</sub>] (2) as starting materials. We undertook this study prompted by (a) the interesting results obtained with such ligands in ruthenium carbonyl cluster chemistry (mentioned above), (b) the publication of an expeditious preparation of the trinuclear cluster 1 directly from [Re<sub>2</sub>(CO)<sub>8</sub>(MeCN)<sub>2</sub>] and hydrogen (90% isolated yield on a half-gram scale),<sup>18</sup> and (c) the publication of the synthesis of some trinuclear carbonyl derivatives by reaction of the tetranuclear cluster 2 with pyridine<sup>19</sup> and acetonitrile.<sup>20</sup>

### Scheme 1



## Results and Discussion

**Reactions of [Re<sub>3</sub>(μ-H)<sub>3</sub>(CO)<sub>11</sub>(MeCN)] (1).** No reaction was observed between complex 1 and 2-hydroxy-6-methylpyridine (Hhmpy) in toluene at reflux temperature. However, under analogous conditions, 2-amino-6-methylpyridine (Hampy) and 2-mercaptopyridine (Hmerpy) reacted slowly with complex 1 (5–7 h) to give the binuclear derivatives [Re<sub>2</sub>(μ-ampy)<sub>2</sub>(CO)<sub>6</sub>] (3) and [Re<sub>2</sub>(μ-merpy)<sub>2</sub>(CO)<sub>6</sub>] (4) (Scheme 1). At least 3 equiv of the corresponding pyridine ligand was necessary to consume all the starting rhenium material. Complex 3 is a new compound, but Deeming et al. have previously reported the synthesis of complex 4 by reaction of [Re<sub>2</sub>(CO)<sub>10</sub>] with Hmerpy in refluxing xylene, including an X-ray structural characterization.<sup>21a</sup> Adams et al. have reported the structure of the closely related quinoline-2-thiolate derivative [Re<sub>2</sub>(μ-merqui)<sub>2</sub>(CO)<sub>6</sub>] (Hmerqui = 2-mercaptoquinoline).<sup>21b</sup>

The IR spectra of 3 and 4 are very similar in the carbonyl region, suggesting analogous structures. This was confirmed by the NMR spectra of 3, which indicate C<sub>2</sub> symmetry for the compound, since only three carbonyl resonances of equal intensity are observed in the <sup>13</sup>C{<sup>1</sup>H} spectrum and only one set of ampy resonances are observed in both the <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} spectra. As far as we are aware, the behavior of 2-aminopyridinate ligands as five-electron donors bridging two metal atoms has no precedent in the chemical literature. However, for 2-mercaptopyridinate ligands, such behavior has been previously observed in binuclear rhenium,<sup>21a</sup> manganese,<sup>22</sup> and molybdenum<sup>23</sup> complexes and in heteronuclear rhenium–ruthenium compounds.<sup>24</sup>

The lower reactivity of Hhmpy, as compared with that of Hampy and Hmerpy, may be explained in terms of the high contribution of the ketonic pyridin-2(1H)-one tautomer, both in solution<sup>25</sup> and in the solid state.<sup>26</sup> In this tautomer the pyridinic nitrogen bears a proton, and therefore the compound is much less basic than Hampy and Hmerpy.

(6) Adams, R. D.; Cortopassi, J. E.; Falloon, S. B. *Organometallics* **1992**, *11*, 3794.

(7) Beringhelli, T.; Ciani, G.; D'Alfonso, G.; Molinari, H.; Sironi, A. *Inorg. Chem.* **1985**, *24*, 2666.

(8) Churchill, M. R.; Bird, P. H.; Kaesz, H. D.; Bau, R.; Fontal, B. *J. Am. Chem. Soc.* **1968**, *90*, 7135.

(9) Andreu, P. L.; Cabeza, J. A.; Riera, V.; Jeannin, Y.; Miguel, D. *J. Chem. Soc., Dalton Trans.* **1990**, 2201.

(10) Lugan, N.; Laurent, F.; Lavigne, G.; Newcomb, T. P.; Liihatta, E. W.; Bonnet, J. *J. Organometallics* **1992**, *11*, 1351.

(11) Andreu, P. L.; Cabeza, J. A.; Fernández-Colinas, J. M.; Riera, V. *J. Chem. Soc., Dalton Trans.* **1990**, 2927.

(12) Deeming, A. J.; Hardcastle, K. I.; Karim, M. *Inorg. Chem.* **1992**, *31*, 4792.

(13) Shen, J. K.; Basolo, F.; Nombel, P.; Lugan, N.; Lavigne, G. *Inorg. Chem.* **1996**, *35*, 755.

(14) (a) Cabeza, J. A.; del Río, I.; Llamazares, A.; Riera, V.; García-Granda, S.; Van der Maelen, J. F. *Inorg. Chem.* **1995**, *34*, 1620. (b) Cabeza, J. A.; del Río, I.; Riera, V.; García-Granda, S.; Sanni, S. B. *Organometallics* **1997**, *16*, 1743.

(15) (a) Cabeza, J. A.; Fernández-Colinas, J. M.; García-Granda, S.; Llamazares, A.; López-Ortiz, F.; Riera, V.; Van der Maelen, J. F. *Organometallics* **1994**, *13*, 426. (b) Cabeza, J. A.; del Río, I.; Riera, V.; Grepioni, F. *Organometallics* **1995**, *14*, 3124. (c) Cabeza, J. A.; del Río, I.; Riera, V.; Grepioni, F. *Organometallics* **1997**, *16*, 812.

(16) (a) Cabeza, J. A.; Fernández-Colinas, J. M.; Llamazares, A. *Synlett* **1995**, 579. (b) Cabeza, J. A.; del Río, I.; Fernández-Colinas, J. M.; Riera, V. *Organometallics* **1996**, *15*, 449.

(17) Nombel, P.; Lugan, N.; Mulla, F.; Lavigne, G. *Organometallics* **1994**, *13*, 4673.

(18) Bruce, M. I.; Low, P. J. *J. Organomet. Chem.* **1996**, *519*, 221.

(19) Wang, S. R.; Cheng, C. P. *J. Organomet. Chem.* **1995**, *490*, 111.

(20) Beringhelli, T.; D'Alfonso, G.; Zarini, M. *J. Chem. Soc., Dalton Trans.* **1995**, 2407.


(21) (a) Deeming, A. J.; Karim, M.; Bates, P. A.; Hursthouse, M. B. *Polyhedron* **1988**, *7*, 1401. (b) Adams, R. D.; Chen, L.; Wu, W. *Organometallics* **1993**, *12*, 4962.

(22) Kabir, S. E.; Karim, M.; Kundu, K.; Ullah, S. M. B.; Hardcastle, K. I. *J. Organomet. Chem.* **1996**, *517*, 155.


(23) (a) Shi, Y.; Lu, S.; Guo, H.; Wu, Q.; Hu, N. *J. Organomet. Chem.* **1996**, *514*, 183. (b) Yu, P.; Huang, L.; Zhuang, B. *Acta Crystallogr.* **1994**, *C50*, 1191.

(24) Cockerton, B.; Deeming, A. J.; Karim, M.; Hardcastle, K. I. *J. Chem. Soc., Dalton Trans.* **1991**, 431.





**Figure 2.** Structure of the anion  $[\text{Re}_3(\mu\text{-H})_3(\mu_3\text{-merpy})(\text{CO})_9]^-$  (**6**) in  $[\text{HNEt}_3]_6$ .



**Figure 3.** Hydrogen-bonding interactions in  $[\text{HNEt}_3]_5$ .


**Table 1. Relevant Bond Lengths (Å) in the Anions  $[\text{Re}_3(\mu\text{-H})_3(\mu_3\text{-ampy})(\text{CO})_9]^-$  (**5**) and  $[\text{Re}_3(\mu\text{-H})_3(\mu_3\text{-merpy})(\text{CO})_9]^-$  (**6**)**

	<b>5</b>	<b>6</b>
Re(1)–Re(2)	3.136(1)	3.113(2)
Re(1)–Re(3)	2.927(1)	2.973(2)
Re(2)–Re(3)	3.154(2)	3.118(2)
Re(1)–C(2)	1.910(8)	1.89(2)
Re(1)–C(1)	1.915(8)	1.91(2)
Re(1)–C(3)	1.906(9)	1.86(2)
Re(3)–C(4)	1.907(9)	1.90(2)
Re(3)–C(5)	1.886(8)	1.96(3)
Re(3)–C(6)	1.928(9)	1.91(2)
Re(2)–C(7)	1.885(8)	1.88(2)
Re(2)–C(8)	1.910(9)	1.93(2)
Re(2)–C(9)	1.924(8)	1.95(2)
mean C–O	1.15(1)	1.16(2)
Re–H <sub>hydride</sub>	1.83	1.83
Re(2)–N(1)	2.250(7)	2.18(1)
Re(1)–N(2)	2.197(6)	
Re(3)–N(2)	2.187(6)	
N(1)–C(10)	1.357(10)	1.35(2)
N(1)–C(14)	1.366(9)	1.33(32)
N(2)–C(10)	1.405(9)	
N(2)–H(2)	1.023(6)	
Re(1)–S		2.450(4)
Re(3)–S		2.453(4)
S–C(10)		1.77(2)
C(10)C(11)	1.38(1)	1.38(2)
C(11)C(12)	1.35(1)	1.36(2)
C(12)C(13)	1.37(1)	1.40(2)
C(13)C(14)	1.37(1)	1.35(2)
C(14)C(15)	1.51(1)	

closely related and will be described together. Relevant structural parameters for both compounds are given in Table 1. The metal frame consists of a Re triangle, with three hydride ligands bridging the three metal–metal bonds. Two hydride ligands are nearly in the metallic

**Table 2. Relevant Hydrogen Bond Interactions in  $[\text{HNEt}_3]_5$  and  $[\text{HNEt}_3]_6$**

N–H...O	N–H (Å)	H...O (Å)	N...O (Å)	N–H...O (deg)
<b><math>[\text{HNEt}_3]_5</math></b>				
N(2)–H(2)...O(2)	1.023	2.522	3.340	136.63
N(2)–H(2)...O(4)	1.023	2.421	3.243	136.86
N(3)–H(3)...O(5)	1.018	1.896	2.889	164.94
<b><math>[\text{HNEt}_3]_6</math></b>				
N(2)–H(2)...O(2)	1.027	2.227	3.071	138.29
N(2)–H(2)...O(6)	1.027	2.437	3.227	131.07



**Figure 4.** Hydrogen-bonding interactions in  $[\text{HNEt}_3]_6$ .

plane, while the plane formed by the remaining hydride ligand and the edge it spans is nearly perpendicular to a Re atom, while the pyridine fragment is attached to a Re atom, while the remaining two Re atoms are bridged by the amido (in **5**) or thiolate (in **6**) group. The Re–Re bond distances range from 2.927(1) to 3.154(2) in **5** and from 2.973(2) to 3.118(2) Å in **6**, the shorter distance being that spanned by the amido (in **5**) or thiolate (in **6**) group. The anions have an idealized  $C_s$  symmetry, with the pseudomirror plane bisecting the Re(1)–Re(3) bond and comprising the remaining Re atom. Nine carbonyl ligands are terminally bound, three to each Re atom. The ampy and merpy ligands contribute five electrons to the clusters, their coordination parameters being comparable to those previously found in triruthenium clusters.<sup>9–11,14–17</sup>

Crystalline  $[\text{HNEt}_3]_5$  presents additional interesting structural features. A short N–H...O hydrogen-bonding interaction is observed between the hydrogen atom of the ammonium cation and one carbonyl ligand of the anion, resulting in the formation of an ionic pair. In addition, two of these pairs are interconnected via N–H...O bifurcated interactions between the N–H group of the ampy ligand and the oxygen atoms of two carbonyl ligands, as shown in Figure 3. The structural parameters of these interactions are listed in Table 2.

The crystal structure of  $[\text{HNEt}_3]_6$  also shows important N–H...O interactions. The basic pattern recognizable in this solid is also a dimer of ionic pairs, in which the H atom of the ammonium cation interacts in a bifurcated way with two carbonyl ligands belonging to two different anionic clusters. This dimeric unit is shown in Figure 4. The relevant structural data of these interactions are listed in Table 2.

The involvement of carbonyl ligands in hydrogen bonding is currently receiving considerable attention in the field of solid-state intermolecular interactions.<sup>32</sup>

### Final Remarks

Probably, the most interesting feature of this article is that it reports convenient syntheses of anionic face-

bridged trinuclear rhenium carbonyl clusters. It is anticipated that reactivity studies on these compounds will lead to interesting derivatives, which will broaden the so far scarce derivative chemistry known for rhenium carbonyl clusters. Such studies are in progress in our laboratories.

### Experimental Section

**General Data.** Solvents were dried over sodium diphenyl ketyl (diethyl ether, hydrocarbons), sodium sulfate (acetone), or calcium hydride (1,2-dichloroethane, dichloromethane) and distilled under nitrogen prior to use. The reactions were carried out under nitrogen, using Schlenk-vacuum line techniques, and were routinely monitored by solution IR spectroscopy (carbonyl stretching region) and by spot TLC (silica gel). Compounds **1**<sup>18</sup> and **2**<sup>3d</sup> were prepared from literature procedures. The pyridines were purchased from Aldrich. Infrared spectra were recorded on a Perkin-Elmer FT 1720-X spectrophotometer, using 0.1 mm CaF<sub>2</sub> cells. <sup>1</sup>H and <sup>13</sup>C NMR spectra were run at room temperature with Bruker AC-200 and AC-300 instruments, being referenced to internal SiMe<sub>4</sub> (for <sup>1</sup>H, δ = 0 ppm) or acetone-*d*<sub>6</sub> (for <sup>13</sup>C, δ = 205.1 ppm). Microanalyses were obtained from the University of Oviedo Analytical Service.

**[Re<sub>2</sub>(μ-ampy)<sub>2</sub>(CO)<sub>6</sub>] (3).** A solution of **1** (50 mg, 0.055 mmol) and Humpy (24 mg, 0.222 mmol) in toluene (10 mL) was stirred at reflux temperature. After complete conversion of the starting material (5 h, IR monitoring), the reaction mixture was cooled to room temperature, leading to precipitation of a white solid. After decantation of the liquid phase, the solid was recrystallized from dichloromethane–hexane to give compound **3** as colorless crystals (35 mg, 56% based on Re). Anal. Found: C, 28.26; H, 1.52; N, 7.62. Calcd for C<sub>18</sub>H<sub>14</sub>N<sub>4</sub>O<sub>6</sub>Re<sub>2</sub>: C, 28.65; H, 1.87; N, 7.43. IR (CH<sub>2</sub>Cl<sub>2</sub>): 2026 (s), 2009 (s), 1916 (sh), 1897 (s), cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.46 (t, *J* = 7.7 Hz, 1 H), 6.53 (d, 7.7 Hz, 1 H), 6.41 (d, 7.7 Hz, 1 H), 3.09 (s, br, 1 H, NH), 2.07 (s, 3 H, Me). <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ(CO) 198.8, 197.1, 194.1; δ(ampy) 173.8, 155.1, 139.3, 117.2, 114.0, 29.6 ppm.

**[Re<sub>2</sub>(μ-merpy)<sub>2</sub>(CO)<sub>6</sub>] (4).** A solution of **1** (50 mg, 0.055 mmol) and Humpy (25 mg, 0.225 mmol) in toluene (10 mL) was stirred at reflux temperature with constant IR monitoring. After complete conversion of the starting material (7 h), the solvent was removed under reduced pressure and the residue was recrystallized from dichloromethane–hexane to give compound **4** as colorless crystals (53 mg, 85% based on Re). Its analytical (CHN%) and spectroscopic data (IR, <sup>1</sup>H NMR) matched those reported in the literature.<sup>21</sup>

**[H<sub>2</sub>ampy][Re<sub>3</sub>(μ-H)<sub>3</sub>(μ<sub>3</sub>-ampy)(CO)<sub>9</sub>] (H<sub>2</sub>ampy)5).** A solution of Humpy (18 mg, 0.160 mmol) in 1,2-dichloroethane (5 mL) was added to a solution of compound **2** (65 mg, 0.060 mmol) in the same solvent (5 mL). After 5 min, the color changed from deep-red to yellow. The reaction mixture was then heated at reflux temperature for 10 min. On cooling to room temperature, a small amount of a white solid precipitated. The solid was filtered off and was subsequently found to be insoluble in dichloromethane and acetone. The filtered solution was evaporated to dryness to give a pale yellow oil, which, upon recrystallization from acetone–hexane, afforded [H<sub>2</sub>ampy]5 as an off-white solid (47 mg, 77% based on Re). Anal. Found: C, 24.31; H, 2.05; N, 5.58. Calcd for C<sub>21</sub>H<sub>19</sub>N<sub>4</sub>O<sub>9</sub>Re<sub>3</sub>: C, 24.49; H, 1.86; N, 5.44. IR (CH<sub>2</sub>Cl<sub>2</sub>): 2029 (s), 2001 (vs), 1911 (vs, br), cm<sup>-1</sup>. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ 7.78–6.66

**Table 3. Crystallographic and Refinement Data for [HNET<sub>3</sub>]5 and [HNET<sub>3</sub>]6**

	[HNET <sub>3</sub> ]5	[HNET <sub>3</sub> ]6
formula	C <sub>21</sub> H <sub>26</sub> N <sub>3</sub> O <sub>9</sub> Re <sub>3</sub>	C <sub>20</sub> H <sub>23</sub> N <sub>2</sub> O <sub>9</sub> Re <sub>3</sub> S
fw	1023.05	1026.06
cryst syst	triclinic	triclinic
space group	<i>P</i> 1	<i>P</i> 1
<i>a</i> , Å	9.479(3)	9.173(6)
<i>b</i> , Å	11.103(3)	11.477(7)
<i>c</i> , Å	13.891(7)	12.922(6)
α, deg	83.51(4)	90.41(4)
β, deg	70.92(4)	91.89(4)
γ, deg	84.78(3)	105.96(5)
volume, Å <sup>3</sup>	1370.5(9)	1307.1(13)
<i>Z</i>	2	2
<i>F</i> (000)	940	940
<i>D</i> <sub>calcd</sub> , g/cm <sup>3</sup>	2.479	2.607
μ, mm <sup>-1</sup>	13.266	13.987
radiation (λ, Å)	Mo Kα (0.710 69)	Mo Kα (0.710 69)
monochromator	graphite	graphite
cryst size, mm	0.20 × 0.16 × 0.15	0.18 × 0.16 × 0.16
temp, K	293(2)	293(2)
θ limits, deg	3.0–27.0	3.0–25.0
min/max <i>h</i> , <i>k</i> , <i>l</i>	–11/12, –13/14, 0/17	–10/10, –13/13, 0/15
no. of reflns collected	5768	4305
no. of unique reflns	5758	2244
no. of reflns with <i>I</i> > 2σ( <i>I</i> )	4851	2122
no. of parameters	325	317
GOF (on <i>F</i> <sup>2</sup> )	0.994	1.007
R1 (on <i>F</i> , <i>I</i> > 2σ( <i>I</i> ))	0.0371	0.0544
wR2 (on <i>F</i> <sup>2</sup> , all data)	0.1025	0.1403

(complex mixture of signals), 5.68 (s, br, 1 H, NH), 2.70 (s, 3 H, Me), 2.49 (s, 3 H, Me), –10.04 (s, 1 H, μ-H), –10.29 (s, 2 H, μ-H) ppm.

**[HNET<sub>3</sub>][Re<sub>3</sub>(μ-H)<sub>3</sub>(μ<sub>3</sub>-ampy)(CO)<sub>9</sub>] (HNET<sub>3</sub>)5).** A solution of Humpy (8 mg, 0.074 mmol) and triethylamine (10 μL, 0.074 mmol) in 1,2-dichloroethane (5 mL) was added to a solution of compound **2** (60 mg, 0.055 mmol) in the same solvent (5 mL). The resulting solution was worked up as described above for [H<sub>2</sub>ampy]5, to give [HNET<sub>3</sub>]5 as a white solid (40 mg, 70%). Anal. Found: C, 25.07; H, 2.61; N, 3.88. Calcd for C<sub>21</sub>H<sub>26</sub>N<sub>3</sub>O<sub>9</sub>Re<sub>3</sub>: C, 24.66; H, 2.54; N, 4.10. IR (CH<sub>2</sub>Cl<sub>2</sub>): 2029 (s), 2001 (vs), 1911 (vs, br), cm<sup>-1</sup>. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ 7.34 (t, *J* = 7.2 Hz, 1 H), 6.87 (d, *J* = 7.2 Hz, 1 H), 6.62 (d, *J* = 7.2 Hz, 1 H), 5.57 (s, br, 1 H, NH), 3.50 (q, *J* = 7.2 Hz, 6 H, CH<sub>2</sub> of Et), 2.69 (s, 3 H, Me), 1.42 (t, *J* = 7.2 Hz, 9 H, CH<sub>3</sub> of Et), –10.08 (s, 1 H, μ-H), –10.31 (s, 2 H, μ-H) ppm. <sup>13</sup>C{<sup>1</sup>H} NMR (acetone-*d*<sub>6</sub>): δ(CO) 195.9 (2 C), 193.5 (1 C), 193.1 (2 C), 188.8 (2 C), 186.1 (2 C); δ(ampy) 181.9, 158.4, 136.9, 116.5, 113.0, 30.1; δ([HNET<sub>3</sub>]<sup>+</sup>) 46.6, 8.3 ppm.

**Reaction of Compound 2 with Hmerpy.** A solution of Hmerpy (18 mg, 0.162 mmol) in 1,2-dichloroethane (5 mL) was added to a solution of compound **2** (60 mg, 0.055 mmol) in the same solvent (5 mL). After 5 min, the color changed from deep-red to yellow. The reaction mixture was then heated at reflux temperature for 10 min. The solution was evaporated to dryness to give an off-white solid. The <sup>1</sup>H NMR spectrum of this material indicated a ca. 3:1 mixture of the binuclear compound **4** and the trinuclear [H<sub>2</sub>merpy]6 (by comparison with the spectra of pure **4** and [HNET<sub>3</sub>]6), but the mixture could not be separated.

**[HNET<sub>3</sub>][Re<sub>3</sub>(μ-H)<sub>3</sub>(μ<sub>3</sub>-merpy)(CO)<sub>9</sub>] (HNET<sub>3</sub>)6).** A solution of Hmerpy (9 mg, 0.081 mmol) and triethylamine (11 μL, 0.08 mmol) in 1,2-dichloroethane (5 mL) was added to a solution of compound **2** (60 mg, 0.055 mmol) in the same solvent (5 mL). The color changed from deep-red to yellow. The solution was heated at reflux temperature for 20 min, and then it was evaporated to dryness. The pale-yellow residue was washed with a small amount of dichloromethane (2 × 1 mL), leaving a white solid. The solid was recrystallized from acetone–hexane to give [HNET<sub>3</sub>]6 as a white solid (31 mg, 55%). Anal. Found: C, 22.93; H, 2.08; N, 2.48. Calcd for C<sub>20</sub>H<sub>23</sub>N<sub>2</sub>O<sub>9</sub>Re<sub>3</sub>S: C, 23.41; H, 2.26; N, 2.73. IR (CH<sub>2</sub>Cl<sub>2</sub>): 2035 (s), 2004 (vs), 1917 (vs, br), 1904 (sh) cm<sup>-1</sup>. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ 8.83 (d, *J* = 7.5 Hz, 1 H), 7.76 (d, *J* = 7.5 Hz, 1 H), 7.63

(32) (a) Grepioni, F.; Braga, D. *Acc. Chem. Res.* **1997**, *30*, 81. (b) Braga, D.; Grepioni, F.; Tedesco, E.; Biradha, K.; Desiraju, G. R. *Organometallics* **1997**, *16*, 1846. (c) Braga, D.; Grepioni, F.; Tedesco, E.; Biradha, K.; Desiraju, G. R. *Organometallics* **1996**, *15*, 2692. (d) Braga, D.; Grepioni, F.; Biradha, K.; Pedireddi, V. R.; Desiraju, G. R. *J. Am. Chem. Soc.* **1995**, *117*, 3156.

(t,  $J = 7.5$  Hz, 1 H), 7.10 (t,  $J = 7.7$  Hz, 1 H), 3.48 (q,  $J = 7.2$  Hz, 6 H,  $\text{CH}_2$  of Et), 1.41 (t,  $J = 7.2$  Hz, 9 H,  $\text{CH}_3$  of Et),  $-10.87$  (s, 2 H,  $\mu\text{-H}$ ),  $-11.58$  (s, 1 H,  $\mu\text{-H}$ ) ppm.  $^{13}\text{C}\{^1\text{H}\}$  NMR (acetone- $d_6$ ):  $\delta(\text{CO})$  194.7 (2 C), 193.5 (1 C), 191.7 (2 C), 187.5 (2 C), 186.9 (2 C);  $\delta(\text{merpy})$  175.6, 165.3, 136.0, 129.9, 122.2;  $\delta([\text{HNEt}_3]^+)$  47.2, 8.3 ppm.

**Crystal Structures of Compounds [HNEt<sub>3</sub>]5 and [HNEt<sub>3</sub>]6.** The X-ray diffraction data collections were carried out on a Nonius CAD-4 diffractometer. Crystal data and details of measurements for compounds [HNEt<sub>3</sub>]5 and [HNEt<sub>3</sub>]6 are summarized in Table 3.

The structures were solved by direct methods<sup>33a</sup> followed by difference Fourier syntheses and full-matrix least-squares refinement on  $F^2$  using SHELXL92.<sup>33b</sup> All non-H atoms were treated anisotropically. Absorption correction was applied in both cases by azimuthal scanning of high  $\chi$  reflections (min.–max. absorption correction 0.56–1.00 and 0.49–1.00 for

[HNEt<sub>3</sub>]5 and [HNEt<sub>3</sub>]6, respectively). H atoms bound to C atoms in the ampy and merpy ligands and in the [HNEt<sub>3</sub>]<sup>+</sup> cations were added in calculated positions and were refined riding on their respective C atoms. The H atoms bound to N atoms were directly located from the Fourier maps and were not refined. The positions of the hydride H atoms were calculated using the program XHYDEX,<sup>33c</sup> and their coordinates were included in the Fourier calculations but were not refined. The approximate location of the hydride H atoms was also detectable by visual inspection of space-filling diagrams of the two anions, since suitable niches were invariably found corresponding to the calculated positions. The program SCHAKAL97<sup>33d</sup> was used for all graphical representations.

**Acknowledgment.** This work has been supported by the Spanish DGES (Grant PB95-1042) and the Italian CNR.

**Supporting Information Available:** ORTEP pictures and tables of bond distances and angles, fractional atomic coordinates, and anisotropic thermal parameters for compounds [HNEt<sub>3</sub>]5 and [HNEt<sub>3</sub>]6 (18 pages). Ordering information is given on any current masthead page.

OM980554Z

(33) (a) Sheldrick, G. M. *Acta Crystallogr.* **1990**, *A46*, 467. (b) Sheldrick, G. M. *SHELXL92 Program for Crystal Structure Determination*; University of Göttingen: Göttingen, Germany, 1993. (c) Orpen, A. G. *XHYDEX, a Program for Locating Hydrides*; Bristol University: Bristol, U.K., 1980. (d) Keller, E. *SCHAKAL97 Graphical Representation of Molecular Models*; University of Freiburg: Freiburg, Germany, 1997.